LAPLACE INVARIANTS OF TODA LATTICES WITH THE EXCEPTIONAL CARTAN MATRICES

A.M. Guryeva* **and A.V. Zhiber**†

arXiv:nlin/0512001v1 [nlin.SI] 1 Dec 2005

We show that Toda lattices with the exceptional Cartan matrices $\mathcal{G}_2$, $\mathcal{F}_4$ and $\mathcal{E}_6 - \mathcal{E}_8$ are Liouville type systems. For these systems of equations, we obtain explicit formulas for the invariants and generalized Laplace invariants.

Keywords: Cartan matrices, integrals, Laplace invariants, generalized invariants

Introduction

It is well known that Laplace invariants and transformations of the scalar hyperbolic equations were the object of classical research. Recently the interest to transformations and Laplace invariants was created. It is explained by the existence of a close relation, discovered in a number of works (see [1, 2, 6, 10-14]), between some important properties of nonlinear equations (such as exact integrability and the existence of higher symmetries, conservation laws, and differential substitutions) and the properties of the chain of invariants of the linear equation. One of the most famous examples of explicitly integrable equations in partial derivatives is the Liouville equation

$$u_{xy} = e^{u}. \tag{1}$$

There are different notions of integrability of nonlinear hyperbolic equations. In particular, the definition of the class of exactly integrable nonlinear hyperbolic equations in [3, 10, 13] was based on the finiteness of the chain of Laplace invariants of the linearized equation (this equations called equations of the Liouville type).

The natural next step in such investigations is to generalize these results to systems of equations. In [14], the definition of equations of the Liouville type was extended to nonlinear hyperbolic systems of form

$$u^i_{xy} = F^i(x, y, u, u_x, u_y) \tag{2}$$

(hereafter, we assume u and F to be n-dimensional vectors).

For completeness of the presentation in what follows, we recall the main results in [14] pertaining to systems of differential equations. The case of systems of differential equations involves a major problem related to the definition of Laplace invariants. The linearized systems of equations for systems (2) has the form

$$\left(D\overline{D} + aD + b\overline{D} + c\right) v = 0, \tag{3}$$

where D and $\overline{D}$ are the respective total derivative operators with respect to x and y and where a, b and c are the matrices

$$a = -\left(\frac{\partial F^i}{\partial u^j_x}\right), \qquad b = -\left(\frac{\partial F^i}{\partial u^j_y}\right), \qquad c = -\left(\frac{\partial F^i}{\partial u^j}\right).$$

A straightforward generalization of the notion of invariants to the matrix case is as follows. The principal Laplace invariants are defined by the formulas

$$H_1 = D(a) + ba - c \qquad \textit{and} \qquad K_1 = \overline{D}(b) + ab - c, \tag{4}$$

---

*Ufa State Aviation Technology University, Ufa, Russia.

†Institute of Mathematics, Ufa Science Center, RAS, Ufa, Russia, e-mail: zhiber@imat.rb.ru.

and the matrices H_i with $i > 1$ are to be found consecutively from the system of equations

$$\overline{D}(H_i) + a_i H_i - H_i a_{i-1} = 0, \tag{5}$$

$$H_{i+1} = D(a_i) + [b, a_i] - \overline{D}(b) + H_i, \qquad i = 1, 2, \dots, \tag{6}$$

where $a_0 = a$. If H_i for $i \leq m$ and a_i for $i \leq m-1$ are already known, then a_m is determined from Eq. (5), and H_{m+1} is determined from Eq. (6). But if $\det H_m = 0$, then a_m either does not exist in general or is determined up to the kernels the matrices H_m. The choice of the element in the kernel essentially affects the existence of and the explicit formulas for the subsequent invariants.

The elements K_i are determined similarly,

$$D(K_i) + b_i K_i - K_i b_{i-1} = 0, \tag{7}$$

$$K_{i+1} = \overline{D}(b_i) + [a, b_i] - D(a) + K_i, \qquad i = 1, 2, \dots, \tag{8}$$

where $b_0 = b$. We thus face the problem of consistently defining the chain of invariants.

We note that the interesting systems (2) are precisely those with degenerate matrices H_i and K_i.

We define the elements H_i and K_i somewhat differently: if the matrices $H_1, H_2, \dots, H_m$ are known and the equation

$$\overline{D}(X_m) + a_m X_m - X_m a = 0, \qquad X_m = H_m \cdot H_{m-1} \cdots H_1, \tag{9}$$

has a solution a_m, we set

$$H_{m+1} = D(a_m) + [b, a_m] - \overline{D}(b) + H_m, \qquad m = 1, 2, \dots. \tag{10}$$

Similarly, if the elements $K_1, K_2, \dots, K_m$ are already found and there exists a solution b_m of the equation

$$D(Y_m) + b_m Y_m - Y_m b = 0, \qquad Y_m = K_m \cdot K_{m-1} \cdots K_1, \tag{11}$$

we define K_{m+1} as

$$K_{m+1} = \overline{D}(b_m) + [a, b_m] - D(a) + K_m, \qquad m = 1, 2 \dots \ . \tag{12}$$

It is clear that under the solvability condition for Eqs. (9) and (11), formulas (4) and (9) – (12) determine a sequence of matrices H_i and K_i, $i = 1, 2, \dots$.

We note that if relations (5) are satisfied for $i = 1, 2, \dots k$, then Eqs. (9) hold for $m = 1, 2, \dots, k$. The converse statement is not true in general.

By analogy with the scalar case, we call the matrices H_i and K_i defined by formulas (4) and (9) – (12) the Laplace invariants and also call X_i and Y_i, $i = 1, 2, \dots$, the generalized invariants of linearized systems of equations (3).

The conditions for the existence of solutions a_m and b_m of systems of equations (9) and (11) are given in the following lemma.

Lemma . *Systems of equations (9) has a solution if and only if the condition*

$$\left(\overline{D} + a\right)(Ker X_m) \subset Ker X_m \tag{13}$$

is satisfied; systems (11) has a solution if and only if the condition

$$(D + b)(Ker Y_m) \subset Ker Y_m \tag{14}$$

is satisfied.

Because the matrices a_i and b_i determined ambiguously, the invariants H_m and K_m depend on the choice of the matrices $a_1, a_2, \dots, a_{m-1}$ and $b_1, b_2, \dots, b_{m-1}$; therefore, the generalized Laplace invariants X_m and Y_m are generally affected by this choice. There thus arises the question under what conditions the sequences $\{X_i\}$ and $\{Y_i\}$ are well-defined. The following theorem answers this question.

Theorem 1. *If*

$$\begin{gathered}\left(D - b^T\right)(Coker X_i) \subset Coker X_i, \qquad i = 1, 2, \dots, m, \\ Coker X_1 \subset Coker X_2 \subset \dots \subset Coker X_m,\end{gathered} \tag{15}$$

then the generalized invariant X_{m+1} is independent of the choice of the matrices $a_1, a_2, \dots, a_m$.

If

$$\begin{gathered}\left(\overline{D} - a^T\right)(Coker Y_i) \subset Coker Y_i, \qquad i = 1, 2, \dots, m, \\ Coker Y_1 \subset Coker Y_2 \subset \dots \subset Coker Y_m,\end{gathered} \tag{16}$$

then the generalized invariant Y_{m+1} is independent of the choice of the matrices $b_1, b_2, \dots, b_m$.

Therefore, precisely the sequences $\{X_i\}$ and $\{Y_i\}$ (rather than $\{H_i\}$ and $\{K_i\}$) are well-defined; their termination is placed in the basic of the definition of systems (2) of the Liouville type.

Definition . *A system of equations (2) is said to be a system of the Liouville type if conditions (13) – (16) are satisfied and there exist $r \geq 1$ and $s \geq 1$ such that $X_r = Y_s = 0$.*

Directly generalizing Liouville equations (1) to systems (2) gives the truncated Toda lattices [7, 8] related to the Cartan matrices of simple Lie algebras. One of the equivalent forms of writing these systems is

$$u^i_{xy} = \sum_{j=1}^{n} a_{ij} \exp(u^j), \qquad i = 1, 2, \dots, n. \tag{17}$$

It is well known that these systems have x and y integrals (see, e.g., [8, 9]). We also note that, as proved in [8], systems of equations (17) has a complete set of x and y integral if and only if the matrix $A = (a_{ij})$ is equivalent to one of the Cartan matrices of a simple Lie algebra.

In this work, we show that Toda lattices (17) with the exceptional Cartan matrices $\mathcal{G}_2$, $\mathcal{F}_4$, $\mathcal{E}_6 - \mathcal{E}_8$ are Liouville type systems in the sense of the definition given above. We obtain explicit formulas for the invariants and generalized Laplace invariants for these systems of equations.

We note that, as showed in [5], systems of equations (17) with Cartan matrices $\mathcal{A}_n$, $\mathcal{B}_n$, $\mathcal{C}_n$, and $\mathcal{D}_n$ are Liouville type systems.

V.V. Sokolov hypothesized that the values of the index k at which the rank of the generalized Laplace invariants X_k drops coincide with the exponents of the corresponding simple Lie algebra and that the number h for which $X_h = 0$, is equal to the Coxeter number.

1. The Cartan matrix $\mathcal{G}_2$

Systems of equations (17) with the Cartan matrix $\mathcal{G}_2$ has the form

$$u^1_{xy} = 2\exp(u^1) - \exp(u^2), \qquad u^2_{xy} = -3\exp(u^1) + 2\exp(u^2). \tag{18}$$

The invariants H_m and generalized invariants X_m of the linearized system (18), determined from formulas (4), (9), and (10), are evaluated as

$$\begin{aligned}
&X_1 = H_1 = \mathcal{G}_2 S_1, && X_m = \mathcal{G}_2 P^{-1} S_m Q, \quad m = 2, 3, 4, 5, \qquad X_6 = 0,\\
&H_2 = \mathcal{G}_2 P^{-1} Z_2 \mathcal{G}_2^{-1}, && H_m = \left(\mathcal{G}_2 P^{-1} Z_m + Q_m\right) P \mathcal{G}_2^{-1}, \quad m = 3, 4, 5, 6,\\
&Q_m = Q_{m-1} + D(B_{m-1}) + \mathcal{G}_2 P^{-1} D_m, && m = 3, 4, 5, 6, \qquad Q_2 = 0,
\end{aligned}$$

where

$$P = \begin{pmatrix} 3 & 1 \\ 0 & \dfrac{1}{3} \end{pmatrix}, \quad Q = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}, \quad Z_2 = \begin{pmatrix} 0 & 0 \\ -\exp(u^2) & \exp(u^1) \end{pmatrix}, \quad B_m = \begin{pmatrix} b^m_{11} & 0 \\ b^m_{21} & 0 \end{pmatrix},$$

b^m_{11} and b^m_{21}, $m = 2, 3, 4, 5$, are arbitrary elements, and $D_3 = \begin{pmatrix} 0 & 0 \\ -\frac{1}{3}e^{u^2} & 0 \end{pmatrix}$, $D_m = 0,\ \ m = 4, 5, 6$, and

$$\begin{aligned}
&S_1 = \mathrm{diag}(\exp(u^1), \exp(u^2)), && S_2 = \mathrm{diag}(0, \exp(u^1 + u^2)), \\
&S_3 = \mathrm{diag}(0, 4\exp(2u^1 + u^2)), && S_4 = \mathrm{diag}(0, 12\exp(3u^1 + u^2)), \\
&S_5 = \mathrm{diag}(0, 12\exp(3u^1 + 2u^2)), && Z_3 = \mathrm{diag}(0, 4\exp(u^1)), \\
&Z_4 = \mathrm{diag}(0, 3\exp(u^1)), && Z_5 = \mathrm{diag}(0, \exp(u^2)), \quad Z_6 = 0.
\end{aligned}$$

Solutions a_m of Eqs. (9) are then given by

$$a_1 = -\mathcal{G}_2 R_1 \mathcal{G}_2^{-1}, \quad a_m = \left[-\mathcal{G}_2 P^{-1} R_m + B_m\right] P \mathcal{G}_2^{-1}, \quad m = 2, 3, 4, 5,$$

where

$$\begin{aligned}
&R_1 = \mathrm{diag}(u^1_y, u^2_y), && R_2 = \mathrm{diag}(0, u^1_y + u^2_y), && R_3 = \mathrm{diag}(0, 2u^1_y + u^2_y), \\
&R_4 = \mathrm{diag}(0, 3u^1_y + u^2_y), && R_5 = \mathrm{diag}(0, 3u^1_y + 2u^2_y).
\end{aligned}$$

We note that the values of the index m at which the rank of the generalized invariants X_m drops coincide with the exponents 1, 5 of the $\mathcal{G}_2$ systems, and the number $m = 6$ for which $X_m = 0$ is equal to the Coxeter number [4].

2. The Cartan matrix $\mathcal{F}_4$

Systems of equations (17) with the Cartan matrix $\mathcal{F}_4$ has the form

$$\begin{aligned}
u^1_{xy} &= 2\exp(u^1) - \exp(u^2), \\
u^2_{xy} &= -\exp(u^1) + 2\exp(u^2) - \exp(u^3), \\
u^3_{xy} &= -2\exp(u^2) + 2\exp(u^3) - \exp(u^4), \\
u^4_{xy} &= -\exp(u^3) + 2\exp(u^4).
\end{aligned} \tag{19}$$

We write this systems in the matrix form

$$D\overline{D}u = \mathcal{F}_4 U c,$$

where $u = (u^1, u^2, u^3, u^4)^T$ is the column of the unknowns, $c = (1, 1, 1, 1)^T$, and $U = \mathrm{diag}(\exp(u^1), \exp(u^2), \exp(u^3), \exp(u^4))$. The linearization of Eqs. (19) is then

$$D\overline{D}v = \mathcal{F}_4 U v, \qquad v = (v^1, v^2, v^3, v^4)^T. \tag{20}$$

To describe the Laplace invariants and the generalized invariants, we introduce matrices.

The matrix J_k is the upper-triangular matrix of size k all of whose elements in the main diagonal and above it are equal to unity. The matrix E_k is the unit matrix of size k.

The matrix M_k is block matrix: $M_k = \begin{pmatrix} E_{5-k} & 0 \\ 0 & J_{k-1} \end{pmatrix}, \quad k = 2, 3, 4.$

We set $\widetilde{J} = \begin{pmatrix} 2 & 2 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$

The matrices $B_m, \ m = 1, 2, \ldots, 11$ of size 4, are defined as
– the first column of the matrices $B_m, \ m \leq 5$ consist of arbitrary elements,
– the first and second columns of the matrices $B_m, \ m = 6, 7$ consist of arbitrary elements,
– for $m \geq 8$ in additional to the first and second columns, the third column B_m is also arbitrary, with the other element equal to zero.
The diagonal matrices S_m and R_m are given by

$$S_1 = \mathrm{diag}\left(\tfrac{1}{2}\exp(u^1), \tfrac{1}{2}\exp(u^2), \exp(u^3), \exp(u^4)\right),$$
$$S_2 = \mathrm{diag}\left(0, 2\exp(u^1 + u^2), 2\exp(u^2 + u^3), \exp(u^3 + u^4)\right),$$
$$S_3 = \mathrm{diag}\left(0, 4\exp(2u^2 + u^3), 2\exp(u^1 + u^2 + u^3), 2\exp(u^2 + u^3 + u^4)\right),$$
$$S_4 = \mathrm{diag}\left(0, 18\exp(u^1 + 2u^2 + u^3), 4\exp(2u^2 + u^3 + u^4), 2\exp(u^1 + u^2 + u^3 + u^4)\right),$$
$$S_5 = \mathrm{diag}\left(0, 36\exp(2u^1 + 2u^2 + u^3), 18\exp(u^1 + 2u^2 + u^3 + u^4), 4\exp(2u^2 + 2u^3 + u^4)\right),$$
$$S_6 = \mathrm{diag}\left(0, 0, 36\exp(2u^1 + 2u^2 + u^3 + u^4), 18\exp(u^1 + 2u^2 + 2u^3 + u^4)\right),$$
$$S_7 = \mathrm{diag}\left(0, 0, 36\exp(2u^1 + 2u^2 + 2u^3 + u^4), 18\exp(u^1 + 3u^2 + 2u^3 + u^4)\right),$$
$$S_8 = \mathrm{diag}\left(0, 0, 0, 18\exp(2u^1 + 3u^2 + 2u^3 + u^4)\right),$$
$$S_9 = \mathrm{diag}\left(0, 0, 0, 36\exp(2u^1 + 4u^2 + 2u^3 + u^4)\right),$$
$$S_{10} = \mathrm{diag}\left(0, 0, 0, 36\exp(2u^1 + 4u^2 + 3u^3 + u^4)\right),$$
$$S_{11} = \mathrm{diag}\left(0, 0, 0, 36\exp(2u^1 + 4u^2 + 3u^3 + 2u^4)\right), \quad S_{12} = 0,$$
$$R_1 = \mathrm{diag}\left(u^1_y, u^2_y, u^3_y, u^4_y\right),$$
$$R_2 = \mathrm{diag}\left(0, u^1_y + u^2_y, u^2_y + u^3_y, u^3_y + u^4_y\right),$$
$$R_3 = \mathrm{diag}\left(0, 2u^2_y + u^3_y, u^1_y + u^2_y + u^3_y, u^2_y + u^3_y + u^4_y\right),$$
$$R_4 = \mathrm{diag}\left(0, u^1_y + 2u^2_y + u^3_y, 2u^2_y + u^3_y + u^4_y, u^1_y + u^2_y + u^3_y + u^4_y\right),$$
$$R_5 = \mathrm{diag}\left(0, 2u^1_y + 2u^2_y + u^3_y, u^1_y + 2u^2_y + u^3_y + u^4_y, 2u^2_y + 2u^3_y + u^4_y\right),$$
$$R_6 = \mathrm{diag}\left(0, 0, 2u^1_y + 2u^2_y + u^3_y + u^4_y, u^1_y + 2u^2_y + 2u^3_y + u^4_y\right),$$
$$R_7 = \mathrm{diag}\left(0, 0, 2u^1_y + 2u^2_y + 2u^3_y + u^4_y, u^1_y + 3u^2_y + 2u^3_y + u^4_y\right),$$
$$R_8 = \mathrm{diag}\left(0, 0, 0, 2u^1_y + 3u^2_y + 2u^3_y + u^4_y\right),$$
$$R_9 = \mathrm{diag}\left(0, 0, 0, 2u^1_y + 4u^2_y + 2u^3_y + u^4_y\right),$$
$$R_{10} = \mathrm{diag}\left(0, 0, 0, 2u^1_y + 4u^2_y + 3u^3_y + u^4_y\right),$$
$$R_{11} = \mathrm{diag}\left(0, 0, 0, 2u^1_y + 4u^2_y + 3u^3_y + 2u^4_y\right).$$

The matrices Z_m, $m = 1, 2, \ldots, 12$, are given by $Z_1 = U$,

$$Z_2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ -2e^{u^2} & 2e^{u^1} & 0 & 0 \\ 0 & -2e^{u^3} & 2e^{u^2} & 0 \\ 0 & 0 & -e^{u^4} & e^{u^3} \end{pmatrix}, \qquad Z_3 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 2e^{u^2} & 0 \\ 0 & -e^{u^3} & e^{u^1} & 0 \\ 0 & 0 & -e^{u^4} & 2e^{u^2} \end{pmatrix},$$

$$Z_4 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 3e^{u^1} & 3e^{u^2} & 0 \\ 0 & -e^{u^4} & 0 & 2e^{u^2} \\ 0 & 0 & -e^{u^4} & e^{u^1} \end{pmatrix}, \qquad Z_5 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 2e^{u^1} & 0 & 0 \\ 0 & -e^{u^4} & 3e^{u^1} & 3e^{u^2} \\ 0 & 0 & -e^{u^3} & 0 \end{pmatrix},$$

$$Z_6 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & -e^{u^4} & 2e^{u^1} & 0 \\ 0 & 0 & -e^{u^3} & 3e^{u^1} \end{pmatrix}, \qquad Z_7 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^3} & -2e^{u^1} \\ 0 & 0 & 0 & e^{u^2} \end{pmatrix},$$

$$Z_8 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^2} & e^{u^1} \end{pmatrix}, \qquad Z_9 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^2} & 2e^{u^2} \end{pmatrix},$$

$$Z_{10} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^2} & e^{u^3} \end{pmatrix}, \qquad Z_{11} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^2} & e^{u^4} \end{pmatrix},$$

$$Z_{12} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^2} & 0 \end{pmatrix}.$$

The matrices P_m and D_m, $m = 2, 3, \ldots, 12$, are given by

$$P_2 = E_4, \quad P_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_4 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -1/3 & 1 & 0 \\ 0 & 1/3 & -1 & 1 \end{pmatrix},$$

$$P_5 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 3 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & -1 & 3 \end{pmatrix}, \qquad P_6 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & \frac{1}{2} \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 3/2 \end{pmatrix}, \qquad P_7 = E_4,$$

$$P_8 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \qquad P_9 = P_{10} = P_{11} = P_{12} = E_4,$$

$$D_3 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ -e^{u^2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \qquad D_7 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 2e^{u^1} & 0 & 0 \\ 0 & -e^{u^3} & 0 & 0 \end{pmatrix},$$

$$D_k = 0, \quad k = 2, 4, 5, 6, 8, 9, \dots, 12.$$

We have the following theorem.

Theorem 2. *Systems of equations (19) is a Liouville type system. The generalized invariants X_m, the invariants H_m of linearized system (20), determined from formulas (4), (9) and (10), are evaluated as*

$$X_m = \mathcal{F}_4 \widetilde{J}^{-1} A_m S_m A_m^T \left(J_4^{-1}\right)^T, \quad m = 1, 2, \dots, 11, \quad X_{12} = 0,$$

$$H_m = \left[\mathcal{F}_4 \widetilde{J}^{-1} A_m Z_m + Q_{m-1}\right] A_{m-1}^{-1} \widetilde{J} \mathcal{F}_4^{-1}, \qquad m = 1, 2, \dots, 12,$$

where the matrices A_m and Q_{m-1} are evaluated using the recursive relations

$$A_m = A_{m-1} M_k^{-1} P_m^{-1}, \qquad m = 3, 4, \dots, 12,$$

$$k = 4 \;\; npu \;\; m \le 6, \;\; k = 3 \;\; npu \;\; m = 7, \;\; k = 2 \;\; npu \;\; m \ge 8,$$

$$A_0 = \widetilde{J} \mathcal{F}_4^{-1}, \qquad A_1 = \widetilde{J}, \qquad A_2 = E_4,$$

$$Q_{m-1} = \mathcal{F}_4 \widetilde{J}^{-1} A_{m-1} D_m + D(B_{m-1}) + Q_{m-2} A_{m-2}^{-1} A_{m-1},$$

$m = 2, 3, \dots, 12$, $Q_0 = 0$.

The elements a_m are then given by

$$a_m = \left[-\mathcal{F}_4 \widetilde{J}^{-1} A_m R_m + B_m\right] A_m^{-1} \widetilde{J} \mathcal{F}_4^{-1}, \qquad m = 1, 2, \dots, 11.$$

The matrices S_m, R_m, and Z_m were calculated by means of programm «Invariant», intended for Maple V Release 4.

We note that the values of the index m at which the rank of the generalized invariants X_m drops coincide with the exponents 1, 5, 7, 11 of the $\mathcal{F}_4$ systems, and the number $m = 12$ for which $X_m = 0$ is equal to the Coxeter number [4].

3. The Cartan matrices $\mathcal{E}_6 - \mathcal{E}_8$

Systems of equations (17) with the Cartan matrix $\mathcal{E}_\alpha$, $\alpha = 6, 7, 8$ has the form

$$\begin{array}{ll}
u^1_{xy} = 2\exp(u^1) - \exp(u^3), & \\
u^2_{xy} = 2\exp(u^2) - \exp(u^4), & \\
u^3_{xy} = -\exp(u^1) + 2\exp(u^3) - \exp(u^4), & \\
u^4_{xy} = -\exp(u^2) - \exp(u^3) + 2\exp(u^4) - \exp(u^5), & \\
u^i_{xy} = -\exp(u^{i-1}) + 2\exp(u^i) - \exp(u^{i+1}), & i = 5, 6, \dots, \alpha - 1, \\
u^\alpha_{xy} = -\exp(u^{\alpha-1}) + 2\exp(u^\alpha). &
\end{array} \tag{21}$$

The linearization of Eqs. (21) is then

$$D\overline{D}v = \mathcal{E}_\alpha U v. \tag{22}$$

The result in this section can be formulated as follows.

Theorem 3. *Systems of equations (21) is a Liouville type system. The generalized invariants X_m, the invariants H_m of linearized system (22), determined from formulas (4), (9), and (10), are evaluated as*

$$X_m = \mathcal{E}_\alpha A_m S_m A_m^T, \quad m = 1, 2, \ldots, r-1, \quad X_r = 0,$$

$$H_m = \left[\mathcal{E}_\alpha A_m Z_m + Q_{m-1}\right] A_{m-1}^{-1} \mathcal{E}_\alpha^{-1}, \qquad m = 1, 2, \ldots, r,$$

where matrices A_m and Q_{m-1} are evaluated using the recursive relations

$$A_m = A_{m-1} G_m^{-1} P_m^{-1}, \quad m = 2, 4, \ldots, r, \quad A_0 = \mathcal{E}_\alpha^{-1}, \ A_1 = E_\alpha,$$

$$Q_{m-1} = \mathcal{E}_\alpha A_{m-1} D_m + D(B_{m-1}) + Q_{m-2} A_{m-2}^{-1} A_{m-1}, \ \ m = 2, 3, \ldots, r, \ \ Q_0 = 0.$$

The elements a_m are then given by

$$a_m = \left[-\mathcal{E}_\alpha A_m R_m + B_m\right] A_m^{-1} \mathcal{E}_\alpha^{-1}, \qquad m = 1, 2, \ldots, r-1.$$

The number r is equal 12 if α set to 6; r is equal 18 if α set to 7 and r is equal 30 if α set to 8.

The matrices S_m, R_m, and Z_m were calculated by means of programm «Invariant», intended for Maple V Release 4.

The matrices P_m, D_m, G_m B_m, S_m, R_m, Z_m are given by formulaes for each $\alpha = 6, 7, 8$ their.

We recall that J_k is the upper-triangular matrix of size k all of whose elements in the main diagonal and above it are equal to unity, and the matrix E_k is the unit matrix of size k.

To find the Laplace invariants of Eqs. (22) when $\alpha = 6$, we introduce matrices of size 6.

The matrices G_m, $m = 2, 3, \ldots, 12$ are given by $G_2 = J_6$,

$$G_m = \begin{pmatrix} 1 & 0 \\ 0 & J_5 \end{pmatrix}, \quad m = 3, 4, 5; \quad G_6 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & J_4 & 0 \\ 0 & 0 & 1 \end{pmatrix};$$

$$G_m = \begin{pmatrix} 1 & 0 & 0 \\ 0 & J_3 & 0 \\ 0 & 0 & E_2 \end{pmatrix}, \qquad m = 7, 8; \quad G_m = E_6 \ \ for \ \ m = 9, 10, 11, 12.$$

The matrices B_m, $m = 1, 2, \ldots, 11$ of size 6, are defined as

- the first column of the matrices B_m, $m \leq 4$ consist of arbitrary elements,
- for $m = 5$ in additional to the first column, the last column B_m is also arbitrary,
- the first, fifth and sixth columns of the matrices B_m, $m = 6, 7$ consist of arbitrary elements,
- the first, fourth, fifth and sixth columns of the matrices B_8 consist of arbitrary elements,
- for $m \geq 9$ in additional to the first, fourth, fifth and sixth columns, the third column B_m is also arbitrary,

with the other element equal to zero.

The matrices P_m, $m = 2, 3, \ldots, 12$ are given by

$$P_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \qquad P_3 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & -1 & 1 & 0 & 0 \\ 0 & -\frac{1}{2} & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_4 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{3} & 1 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & -1 & 1 & 0 & 0 \\ 0 & -\frac{1}{6} & \frac{1}{2} & -1 & 1 & 0 \\ 0 & \frac{1}{6} & -\frac{1}{2} & 1 & -1 & 1 \end{pmatrix}, \qquad P_5 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & -\frac{1}{3} & 1 & 0 & 0 \\ 0 & -\frac{1}{3} & \frac{1}{3} & -1 & 1 & 0 \\ 0 & \frac{1}{6} & -\frac{1}{6} & \frac{1}{2} & -1 & 1 \end{pmatrix},$$

$$P_6 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{3}{5} & 1 & 0 & 0 & 0 \\ 0 & \frac{3}{5} & -1 & 1 & 0 & 0 \\ 0 & -\frac{1}{5} & \frac{1}{3} & -\frac{1}{3} & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \qquad P_7 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & \frac{5}{3} & 0 & 0 \\ 0 & 0 & 1 & -\frac{5}{3} & 0 & 0 \\ 0 & 0 & \frac{5}{3} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_8 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{5}{7} & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & \frac{2}{7} & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \qquad P_9 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_{10} = P_{11} = P_{12} = E_6.$$

The nonzero elements of the matrices $D_m = (d_{ij})_{i,j=1}^{6}$, $m = 2, 3, \ldots, 12$, are determined by the formulas

$$d_{21}^{3} = -e^{u^3}; \;\; d_{46}^{6} = 2e^{u^2} - 2e^{u^3}; \;\; d_{56}^{6} = e^{u^1}; \;\; d_{35}^{7} = -e^{u^4}; \;\; d_{55}^{7} = 3e^{u^2} - \tfrac{3}{2}e^{u^1};$$
$$d_{24}^{9} = e^{u^5} - \tfrac{5}{2}e^{u^1}; \quad d_{35}^{9} = e^{u^3}; \quad d_{23}^{10} = e^{u^5}.$$

The diagonal matrices S_m and R_m, are given by

$$S_1 = \mathrm{diag}\left(\exp(u^1), \exp(u^2), \exp(u^3), \exp(u^4), \exp(u^5), \exp(u^6)\right),$$
$$S_2 = \mathrm{diag}\left(0, \exp(u^1+u^3), \exp(u^2+u^4), \exp(u^3+u^4), \exp(u^4+u^5), \exp(u^5+u^6)\right),$$
$$S_3 = \mathrm{diag}\left(0, 4\exp(u^2+u^3+u^4), \exp(u^1+u^3+u^4), \exp(u^2+u^4+u^5), \exp(u^3+u^4+u^5),\right.$$
$$\left.\exp(u^4+u^5+u^6)\right), \qquad S_4 = \mathrm{diag}\left(0, 9\exp(u^1+u^2+u^3+u^4), 4\exp(u^2+u^3+u^4+u^5),\right.$$
$$\left.\exp(u^1+u^3+u^4+u^5), \exp(u^2+u^4+u^5+u^6), \exp(u^3+u^4+u^5+u^6)\right),$$
$$S_5 = \mathrm{diag}\left(0, 4\exp(u^2+u^3+2u^4+u^5), 9\exp(u^1+u^2+u^3+u^4+u^5),\right.$$
$$\left.4\exp(u^2+u^3+u^4+u^5+u^6), \exp(u^1+u^3+u^4+u^5+u^6), 0\right),$$
$$S_6 = \mathrm{diag}\left(0, 25\exp(u^1+u^2+u^3+2u^4+u^5), 4\exp(u^2+u^3+2u^4+u^5+u^6),\right.$$
$$\left.9\exp(u^1+u^2+u^3+u^4+u^5+u^6), 0, 0\right),$$
$$S_7 = \mathrm{diag}\left(0, 25\exp(u^1+u^2+2u^3+2u^4+u^5), 4\exp(u^1+u^2+u^3+2u^4+\right.$$
$$\left.+2u^5+u^6), 25\exp(u^2+u^3+2u^4+u^5+u^6), 0, 0\right),$$

$$
\begin{aligned}
&S_8 = \operatorname{diag}\left(0, 25\exp(u^1+u^2+u^3+2u^4+2u^5+u^6),\right. \\
&\left. 25\exp(u^1+u^2+2u^3+2u^4+u^5+u^6), 0, 0, 0\right), \\
&S_9 = \operatorname{diag}\left(0, 25\exp(u^1+u^2+u^3+2u^4+2u^5+u^6), 0, 0, 0, 0\right), \\
&S_{10} = \operatorname{diag}\left(0, 25\exp(u^1+u^2+2u^3+3u^4+2u^5+u^6), 0, 0, 0, 0\right), \\
&S_{11} = \operatorname{diag}\left(0, 25\exp(u^1+2u^2+2u^3+3u^4+2u^5+u^6), 0, 0, 0, 0\right), \quad S_{12} = 0, \\
&R_1 = \operatorname{diag}\left(u^1_y, u^2_y, u^3_y, u^4_y, u^5_y, u^6_y\right), \\
&R_2 = \operatorname{diag}\left(0, u^1_y+u^3_y, u^2_y+u^4_y, u^3_y+u^4_y, u^4_y+u^5_y, u^5_y+u^6_y)\right), \\
&R_3 = \operatorname{diag}\left(0, u^2_y+u^3_y+u^4_y, u^1_y+u^3_y+u^4_y, u^2_y+u^4_y+u^5_y, u^3_y+u^4_y+u^5_y, u^4_y+u^5_y+u^6_y\right), \\
&R_4 = \operatorname{diag}\left(0, u^1_y+u^2_y+u^3_y+u^4_y, u^2_y+u^3_y+u^4_y+u^5_y, u^1_y+u^3_y+u^4_y+u^5_y, u^2_y+u^4_y+u^5_y+u^6_y,\right. \\
&\left. u^3_y+u^4_y+u^5_y+u^6_y\right), \quad R_5 = \operatorname{diag}\left(0, u^2_y+u^3_y+2u^4_y+u^5_y, u^1_y+u^2_y+u^3_y+u^4_y+u^5_y,\right. \\
&\left. u^2_y+u^3_y+u^4_y+u^5_y+u^6_y, u^1_y+u^3_y+u^4_y+u^5_y+u^6_y, 0\right), \\
&R_6 = \operatorname{diag}\left(0, u^1_y+u^2_y+u^3_y+2u^4_y+u^5_y, u^2_y+u^3_y+2u^4_y+u^5_y+u^6_y,\right. \\
&\left. u^1_y+u^2_y+u^3_y+u^4_y+u^5_y+u^6_y, 0, 0\right), \\
&R_7 = \operatorname{diag}\left(0, u^1_y+u^2_y+2u^3_y+2u^4_y+u^5_y, u^1_y+u^2_y+u^3_y+2u^4_y+2u^5_y+u^6_y,\right. \\
&\left. u^2_y+u^3_y+2u^4_y+u^5_y+u^6_y, 0, 0\right), \\
&R_8 = \operatorname{diag}\left(0, u^1_y+u^2_y+u^3_y+2u^4_y+2u^5_y+u^6_y,\right. \\
&\left. u^1_y+u^2_y+u^3_y+2u^4_y+2u^5_y+u^6_y, 0, 0, 0\right), \\
&R_9 = \operatorname{diag}\left(0, u^1_y+u^2_y+2u^3_y+2u^4_y+2u^5_y+u^6_y, 0, 0, 0, 0\right), \\
&R_{10} = \operatorname{diag}\left(0, u^1_y+u^2_y+2u^3_y+3u^4_y+2u^5_y+u^6_y, 0, 0, 0, 0\right), \\
&R_{11} = \operatorname{diag}\left(0, u^1_y+2u^2_y+2u^3_y+3u^4_y+2u^5_y+u^6_y, 0, 0, 0, 0\right).
\end{aligned}
$$

The matrices $Z_m, \;\; m = 1, 2, \ldots, 12$, are given by $\;\; Z_1 = U$,

$$
Z_2 = \begin{pmatrix}
0 & 0 & 0 & 0 & 0 & 0 \\
-e^{u^3} & 0 & e^{u^1} & 0 & 0 & 0 \\
0 & -e^{u^4} & 0 & e^{u^2} & 0 & 0 \\
0 & 0 & -e^{u^4} & e^{u^3} & 0 & 0 \\
0 & 0 & 0 & -e^{u^5} & e^{u^4} & 0 \\
0 & 0 & 0 & 0 & -e^{u^6} & e^{u^5}
\end{pmatrix}, \quad
Z_3 = \begin{pmatrix}
0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 2e^{u^3} & 2e^{u^2} & 0 & 0 \\
0 & -e^{u^4} & 0 & e^{u^1} & 0 & 0 \\
0 & 0 & -e^{u^5} & 0 & e^{u^2} & 0 \\
0 & 0 & 0 & -e^{u^5} & e^{u^3} & 0 \\
0 & 0 & 0 & 0 & -e^{u^6} & e^{u^4}
\end{pmatrix},
$$

$$
Z_4 = \begin{pmatrix}
0 & 0 & 0 & 0 & 0 & 0 \\
0 & \frac{3}{2}e^{u^1} & 3e^{u^2} & 0 & 0 & 0 \\
0 & -e^{u^5} & 0 & 2e^{u^3} & 2e^{u^2} & 0 \\
0 & 0 & -e^{u^5} & 0 & e^{u^1} & 0 \\
0 & 0 & 0 & -e^{u^6} & 0 & e^{u^2} \\
0 & 0 & 0 & 0 & -e^{u^6} & e^{u^3}
\end{pmatrix}, \quad
Z_5 = \begin{pmatrix}
0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & e^{u^4} & 0 & 0 & 0 \\
0 & -e^{u^5} & \frac{3}{2}e^{u^1} & 3e^{u^2} & 0 & 0 \\
0 & 0 & -e^{u^6} & 0 & 2e^{u^3} & 2e^{u^2} \\
0 & 0 & 0 & -e^{u^6} & 0 & e^{u^1} \\
0 & 0 & 0 & 0 & 0 & 0
\end{pmatrix},
$$

$$Z_6 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{5}{2}e^{u^1} & \frac{5}{3}e^{u^4} & 0 & 0 & 0 \\ 0 & -e^{u^6} & 0 & e^{u^4} & 0 & 0 \\ 0 & 0 & -e^{u^6} & \frac{3}{2}e^{u^1} & 3e^{u^2} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad Z_7 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^3} & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^5} & 0 & 0 & 0 \\ 0 & -e^{u^6} & \frac{5}{2}e^{u^1} & \frac{5}{3}e^{u^4} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_8 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{5}{2}e^{u^1} & e^{u^5} & 0 & 0 \\ 0 & -e^{u^6} & 0 & e^{u^3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad Z_9 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^3} & e^{u^5} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{10} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^4} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad Z_{11} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{12} = 0.$$

To find the Laplace invariants of Eqs. (22) when $\alpha = 7$, we introduce matrices of size 7.
The matrices $B_m,\ \ m = 1, 2, \dots, 17$ of size 7, are defined as

- the first column of the matrices $B_m,\ m \leq 5$ consist of arbitrary elements,
- for $m = 6, 7$ in additional to the first column, the last column B_m is also arbitrary,
- the first, sixth and seventh columns of the matrices $B_m,\ m = 8, 9$ consist of arbitrary elements,
- the first, second, sixth and seventh columns of the matrices $B_m,\ m = 10, 11$ consist of arbitrary elements,
- the first, second, third, sixth and seventh columns of the matrices $B_m,\ m = 12, 13$ consist of arbitrary elements,
- for $m \geq 14$ in additional to the first, second, third, firth, sixth and seventh columns, the third column B_m is also arbitrary,

with the other element equal to zero.

The matrices $P_m,\ \ m = 2, 3, \dots, 12$, are given by

$$P_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_3 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & -1 & 1 & 0 & 0 & 0 \\ 0 & -\frac{1}{2} & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_4 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{3} & 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & -1 & 1 & 0 & 0 & 0 \\ 0 & -\frac{1}{6} & \frac{1}{2} & -1 & 1 & 0 & 0 \\ 0 & \frac{1}{6} & -\frac{1}{2} & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \qquad P_5 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & -\frac{1}{3} & 1 & 0 & 0 & 0 \\ 0 & -\frac{1}{3} & \frac{1}{3} & -1 & 1 & 0 & 0 \\ 0 & \frac{1}{6} & -\frac{1}{6} & \frac{1}{2} & -1 & 1 & 0 \\ 0 & -\frac{1}{6} & \frac{1}{6} & -\frac{1}{2} & 1 & -1 & 1 \end{pmatrix},$$

$$P_6 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{3}{5} & 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{3}{5} & -1 & 1 & 0 & 0 & 0 \\ 0 & -\frac{1}{5} & \frac{1}{3} & -\frac{1}{3} & 1 & 0 & 0 \\ 0 & \frac{1}{5} & -\frac{1}{3} & \frac{1}{3} & -1 & 1 & 0 \\ 0 & -\frac{1}{10} & \frac{1}{6} & -\frac{1}{6} & \frac{1}{2} & -1 & 1 \end{pmatrix}, \qquad P_7 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & \frac{5}{3} & -\frac{5}{3} & 5 & 0 \\ 0 & 0 & 0 & \frac{5}{3} & -\frac{5}{3} & 5 & 0 \\ 0 & 0 & 0 & 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & -1 & \frac{5}{3} & -\frac{5}{3} & 5 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_8 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -6 & 10 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 7 & -10 & 0 & 0 \\ 0 & -\frac{4}{5} & 0 & \frac{9}{5} & -3 & 0 & 0 \\ 0 & -\frac{6}{5} & 0 & \frac{21}{5} & -7 & 1 & 0 \\ 0 & \frac{2}{5} & 0 & -\frac{7}{5} & \frac{7}{3} & -\frac{1}{3} & 1 \end{pmatrix}, \qquad P_9 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 4 & 0 & 0 \\ 0 & -\frac{1}{2} & 1 & 0 & -\frac{1}{2} & 0 & 0 \\ 0 & -\frac{1}{2} & 0 & 1 & -\frac{1}{2} & 0 & 0 \\ 0 & -\frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 1 \end{pmatrix},$$

$$P_{10} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & -2 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -\frac{3}{5} & 1 \end{pmatrix}, \qquad P_{11} = P_{12} = E_7,$$

$$P_{13} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \qquad P_{14} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_{15} = P_{16} = P_{17} = P_{18} = E_7.$$

The diagonal matrices S_m and R_m, are given by

$$S_1 = \mathrm{diag}\left(\exp(u^1), \exp(u^2), \exp(u^3), \exp(u^4), \exp(u^5), \exp(u^6), \exp(u^7)\right),$$

$$S_2 = \mathrm{diag}\left(0, \exp(u^1+u^3), \exp(u^2+u^4), \exp(u^3+u^4), \exp(u^4+u^5), \exp(u^5+u^6), \exp(u^6+u^7)\right),$$

$$S_3 = \mathrm{diag}\left(0, 4\exp(u^2+u^3+u^4), \exp(u^1+u^3+u^4), \exp(u^2+u^4+u^5), \exp(u^4+u^5+u^6),\right.$$

$$\left.\exp(u^5+u^6+u^7)\right), \quad S_4 = \mathrm{diag}\left(0, 9\exp(u^1+u^2+u^3+u^4), 4\exp(u^2+u^3+u^4+u^5),\right.$$

$$\left.\exp(u^1+u^3+u^4+u^5), \exp(u^2+u^4+u^5+u^6), \exp(u^3+u^4+u^5+u^6), \exp(u^4+u^5+u^6+u^7)\right),$$

$$S_5 = \mathrm{diag}\left(0, 4\exp(u^2+u^3+2u^4+u^5), 9\exp(u^1+u^2+u^3+u^4+u^5), 4\exp(u^2+u^3+u^4+u^5+u^6),\right.$$

$$\left.\exp(u^1+u^3+u^4+u^5+u^6), \exp(u^2+u^4+u^5+u^6+u^7), \exp(u^3+u^4+u^5+u^6+u^7)\right),$$

$$S_6 = \mathrm{diag}\left(0, 25\exp(u^1+u^2+u^3+2u^4+u^5), 4\exp(u^2+u^3+2u^4+u^5+u^6),\right.$$

$$9\exp(u^1+u^2+u^3+u^4+u^5+u^6), 4\exp(u^2+u^3+u^4+u^5+u^6+u^7),$$

$$\left.\exp(u^1+u^3+u^4+u^5+u^6+u^7), 0\right), \quad S_7 = \mathrm{diag}\left(0, 25\exp(u^1+u^2+2u^3+2u^4+u^5),\right.$$

$$25\exp(u^1+u^2+u^3+2u^4+u^5+u^6), 4\exp(u^2+u^3+2u^4+u^5+u^6+u^7), 9\exp(u^1+u^2+u^3+u^4+$$
$$\left.u^5+u^6+u^7), 4\exp(u^2+u^3+2u^4+2u^5+u^6+u^7), 0\right),$$

$$S_8 = \mathrm{diag}\left(0, 25\exp(u^1+u^2+u^3+2u^4+2u^5+u^6), 25\exp(u^1+u^2+2u^3+2u^4+u^5+u^6),\right.$$

$$\left.25\exp(u^1+u^2+u^3+2u^4+u^5+u^6+u^7), 4\exp(u^2+u^3+2u^4+2u^5+u^6+u^7), 0, 0\right),$$

$$S_9 = \mathrm{diag}\left(0, 100\exp(u^2+u^3+2u^4+2u^5+2u^6+u^7), 25\exp(u^1+u^2+2u^3+2u^4+2u^5+u^6),\right.$$

$$\left.25\exp(u^1+u^2+2u^3+2u^4+u^5+u^6+u^7), 25\exp(u^1+u^2+u^3+2u^4+2u^5+u^6+u^7), 0, 0\right),$$

$$S_{10} = \mathrm{diag}\left(0, 0, 25\exp(u^1+u^2+2u^3+3u^4+2u^5+u^6), 25\exp(u^1+u^2+2u^3+2u^4+2u^5+u^6+u^7),\right.$$

$$\left.25\exp(u^1+u^2+u^3+2u^4+2u^5+2u^6+u^7), 0, 0\right), \quad S_{11} = \mathrm{diag}\left(0, 0, 0, 25\exp(u^1+2u^2+2u^3+\right.$$

$$+3u^4+2u^5+u^6), 25\exp(u^1+u^2+2u^3+3u^4+2u^5+u^6+u^7), 25\exp(u^1+u^2+2u^3+2u^4+$$

$$\left.+2u^5+2u^6+u^7), 0, 0\right), \quad S_{12} = \mathrm{diag}\left(0, 0, 0, 25\exp(u^1+2u^2+2u^3+3u^4+2u^5+u^6+u^7),\right.$$

$$\left.25\exp(u^1+u^2+2u^3+3u^4+2u^5+2u^6+u^7), 0, 0\right),$$

$$S_{13} = \mathrm{diag}\left(0, 0, 0, 25\exp(u^1+u^2+2u^3+3u^4+3u^5+2u^6+u^7),\right.$$

$$\left.25\exp(u^1+2u^2+2u^3+3u^4+2u^5+2u^6+u^7), 0, 0\right),$$

$$S_{14} = \mathrm{diag}\left(0, 0, 0, 0, 25\exp(u^1+2u^2+2u^3+3u^4+3u^5+2u^6+u^7), 0, 0\right),$$

$$S_{15} = \mathrm{diag}\left(0, 0, 0, 0, 25\exp(u^1+2u^2+2u^3+4u^4+3u^5+2u^6+u^7), 0, 0\right),$$

$$S_{16} = \mathrm{diag}\left(0, 0, 0, 0, 25\exp(u^1+2u^2+3u^3+4u^4+3u^5+2u^6+u^7), 0, 0\right),$$

$$S_{17} = \mathrm{diag}\left(0, 0, 0, 0, 25\exp(2u^1+2u^2+3u^3+4u^4+3u^5+2u^6+u^7), 0, 0\right),$$

$$S_{18} = 0,$$

$$R_1 = \mathrm{diag}\left(u^1_y, u^2_y, u^3_y, u^4_y, u^5_y, u^6_y, u^7_y\right),$$

$$R_2 = \mathrm{diag}\left(0, u^1_y+u^3_y, u^2_y+u^4_y, u^3_y+u^4_y, u^4_y+u^5_y, u^5_y+u^6_y, u^6_y+u^7_y\right),$$

$$R_3 = \mathrm{diag}\left(0, u^2_y+u^3_y+u^4_y, u^1_y+u^3_y+u^4_y, u^2_y+u^4_y+u^5_y, u^4_y+u^5_y+u^6_y, u^5_y+u^6_y+u^7_y\right),$$

$$R_4 = \mathrm{diag}\big(0, u_y^1 + u_y^2 + u_y^3 + u_y^4, u_y^2 + u_y^3 + u_y^4 + u_y^5, u_y^1 + u_y^3 + u_y^4 + u_y^5, u_y^2 + u_y^4 + u_y^5 + u_y^6,$$

$$u_y^3 + u_y^4 + u_y^5 + u_y^6, u_y^4 + u_y^5 + u_y^6 + u_y^7\big)\,,$$

$$R_5 = \mathrm{diag}\big(0, u_y^2 + u_y^3 + 2u_y^4 + u_y^5, u_y^1 + u_y^2 + u_y^3 + u_y^4 + u_y^5, u_y^2 + u_y^3 + u_y^4 + u_y^5 + u_y^6,$$

$$u_y^1 + u_y^3 + u_y^4 + u_y^5 + u_y^6, u_y^2 + u_y^4 + u_y^5 + u_y^6 + u_y^7, u_y^3 + u_y^4 + u_y^5 + u_y^6 + u_y^7\big)\,,$$

$$R_6 = \mathrm{diag}\big(0, u_y^1 + u_y^2 + u_y^3 + 2u_y^4 + u_y^5, u_y^2 + u_y^3 + 2u_y^4 + u_y^5 + u_y^6, u_y^1 + u_y^2 + u_y^3 +$$

$$+ u_y^4 + u_y^5 + u_y^6, u_y^2 + u_y^3 + u_y^4 + u_y^5 + u_y^6 + u_y^7, u_y^1 + u_y^3 + u_y^4 + u_y^5 + u_y^6 + u_y^7, 0\big)\,,$$

$$R_7 = \mathrm{diag}\big(0, u_y^1 + u_y^2 + 2u_y^3 + 2u_y^4 + u_y^5, u_y^1 + u_y^2 + u_y^3 + 2u_y^4 + u_y^5 + u_y^6, u_y^2 + u_y^3 + 2u_y^4 + u_y^5 + u_y^6 + u_y^7,$$

$$u_y^1 + u_y^2 + u_y^3 + u_y^4 + u_y^5 + u_y^6 + u_y^7, u_y^2 + u_y^3 + 2u_y^4 + 2u_y^5 + u_y^6 + u_y^7, 0\big)\,,$$

$$R_8 = \mathrm{diag}\big(0, u_y^1 + u_y^2 + u_y^3 + 2u_y^4 + 2u_y^5 + u_y^6, u_y^1 + u_y^2 + 2u_y^3 + 2u_y^4 + u_y^5 + u_y^6,$$

$$u_y^1 + u_y^2 + u_y^3 + 2u_y^4 + u_y^5 + u_y^6 + u_y^7, u_y^2 + u_y^3 + 2u_y^4 + 2u_y^5 + u_y^6 + u_y^7, 0, 0\big)\,,$$

$$R_9 = \mathrm{diag}\big(0, u_y^2 + u_y^3 + 2u_y^4 + 2u_y^5 + 2u_y^6 + u_y^7, u_y^1 + u_y^2 + 2u_y^3 + 2u_y^4 + 2u_y^5 + u_y^6,$$

$$u_y^1 + u_y^2 + 2u_y^3 + 2u_y^4 + u_y^5 + u_y^6 + u_y^7, u_y^1 + u_y^2 + u_y^3 + 2u_y^4 + 2u_y^5 + u_y^6 + u_y^7, 0, 0\big)\,,$$

$$R_{10} = \mathrm{diag}\big(0, 0, u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + u_y^6, u_y^1 + u_y^2 + 2u_y^3 + 2u_y^4 +$$

$$2u_y^5 + u_y^6 + u_y^7, 25\exp(u^1 + u^2 + u^3 + 2u^4 + 2u^5 + 2u_y^6 + u_y^7, 0, 0\big)\,,$$

$$R_{11} = \mathrm{diag}\big(0, 0, 0, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + u_y^6, u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + u_y^6 + u_y^7,$$

$$u_y^1 + u_y^2 + 2u_y^3 + 2u_y^4 + 2u_y^5 + 2u_y^6 + u_y^7, 0, 0\big)\,,$$

$$R_{12} = \mathrm{diag}\big(0, 0, 0, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + u_y^6 + u_y^7, u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + 2u_y^6 + u_y^7, 0, 0\big)\,,$$

$$R_{13} = \mathrm{diag}\big(0, 0, 0, u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + 2u_y^6 + u_y^7, 0, 0\big)\,,$$

$$R_{14} = \mathrm{diag}\big(0, 0, 0, 0, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7, 0, 0\big)\,,$$

$$R_{15} = \mathrm{diag}\big(0, 0, 0, 0, u_y^1 + 2u_y^2 + 2u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7, 0, 0\big)\,,$$

$$R_{16} = \mathrm{diag}\big(0, 0, 0, 0, u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7, 0, 0\big)\,,$$

$$R_{17} = \mathrm{diag}\big(0, 0, 0, 0, 2u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7, 0, 0\big)\,.$$

The matrices $Z_m,\ \ m = 1, 2, \ldots, 18$, are given by $Z_1 = U$,

$$Z_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -e^{u^3} & 0 & e^{u^1} & 0 & 0 & 0 & 0 \\ 0 & -e^{u^4} & 0 & e^{u^2} & 0 & 0 & 0 \\ 0 & 0 & -e^{u^4} & e^{u^3} & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^5} & e^{u^4} & 0 & 0 \\ 0 & 0 & 0 & 0 & -e^{u^6} & e^{u^5} & 0 \\ 0 & 0 & 0 & 0 & 0 & -e^{u^7} & e^{u^6} \end{pmatrix},$$

$$Z_3 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2e^{u^3} & 2e^{u^2} & 0 & 0 & 0 \\ 0 & -e^{u^4} & 0 & e^{u^1} & 0 & 0 & 0 \\ 0 & 0 & -e^{u^5} & 0 & e^{u^2} & 0 & 0 \\ 0 & 0 & 0 & -e^{u^5} & e^{u^3} & 0 & 0 \\ 0 & 0 & 0 & 0 & -e^{u^6} & e^{u^4} & 0 \\ 0 & 0 & 0 & 0 & 0 & -e^{u^7} & e^{u^5} \end{pmatrix},$$

$$Z_4 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \tfrac{3}{2}e^{u^1} & \tfrac{3}{2}e^{u^2} & 0 & 0 & 0 & 0 \\ 0 & -e^{u^5} & 0 & 2e^{u^3} & 2e^{u^2} & 0 & 0 \\ 0 & 0 & -e^{u^5} & 0 & e^{u^1} & 0 & 0 \\ 0 & 0 & 0 & -e^{u^6} & 0 & e^{u^2} & 0 \\ 0 & 0 & 0 & 0 & -e^{u^6} & e^{u^3} & 0 \\ 0 & 0 & 0 & 0 & 0 & -e^{u^7} & e^{u^4} \end{pmatrix},$$

$$Z_5 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^4} & 0 & 0 & 0 & 0 \\ 0 & -e^{u^5} & \tfrac{3}{2}e^{u^1} & 3e^{u^2} & 0 & 0 & 0 \\ 0 & 0 & -e^{u^6} & 0 & 2e^{u^3} & 2e^{u^2} & 0 \\ 0 & 0 & 0 & -e^{u^6} & 0 & e^{u^1} & 0 \\ 0 & 0 & 0 & 0 & -e^{u^7} & 0 & e^{u^2} \\ 0 & 0 & 0 & 0 & 0 & -e^{u^7} & e^{u^3} \end{pmatrix},$$

$$Z_6 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \tfrac{5}{2}e^{u^1} & \tfrac{5}{3}e^{u^4} & 0 & 0 & 0 & 0 \\ 0 & -e^{u^6} & 0 & e^{u^4} & 0 & 0 & 0 \\ 0 & 0 & -e^{u^6} & \tfrac{3}{2}e^{u^1} & 3e^{u^2} & 0 & 0 \\ 0 & 0 & 0 & -e^{u^7} & 0 & 2e^{u^3} & 2e^{u^2} \\ 0 & 0 & 0 & 0 & -e^{u^7} & 0 & e^{u^1} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_7 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^3} & 0 & 0 & 0 & 0 & 0 \\ 0 & -e^{u^6} & \tfrac{5}{2}e^{u^1} & \tfrac{5}{3}e^{u^4} & 0 & 0 & 0 \\ 0 & 0 & -e^{u^7} & 0 & e^{u^4} & 0 & 0 \\ 0 & 0 & 0 & -e^{u^7} & \tfrac{3}{2}e^{u^1} & 3e^{u^2} & 0 \\ 0 & 0 & -e^{u^5} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_8 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^5} & 0 & 0 & \frac{5}{2}e^{u^1} & 0 \\ 0 & -e^{u^6} & e^{u^3} & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^7} & \frac{5}{2}e^{u^1} & \frac{5}{3}e^{u^4} & 0 & 0 \\ 0 & 0 & 0 & -e^{u^5} & 0 & -e^{u^7} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_9 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 5e^{u^6} & 0 & 0 \\ 0 & -e^{u^3} & e^{u^5} & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^7} & e^{u^3} & 0 & 0 & 0 \\ 0 & -e^{u^7} & 0 & -e^{u^5} & \frac{5}{2}e^{u^1} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{10} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^4} & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^7} & e^{u^5} & -e^{u^3} & 0 & 0 \\ 0 & \frac{1}{2}e^{u^1} & 0 & 0 & e^{u^6} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{11} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^2} & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^7} & e^{u^4} & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^6} & e^{u^3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{12} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^7} & e^{u^2} & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^6} & e^{u^4} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{13} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & e^{u^5} & 0 & 0 \\ 0 & 0 & 0 & -e^{u^6} & e^{u^2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{14} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & e^{u^2} & e^{u^5} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad Z_{15} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & e^{u^4} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{16} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & e^{u^3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad Z_{17} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & e^{u^1} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{18} = 0.$$

The nonzero elements of the matrices $D_m = (d_{ij})_{i,j=1}^{7}$, $m = 2, 3, \ldots, 18$, are determined by the formulas

$$d_{21}^3 = -e^{u^3}; \;\; d_{57}^7 = 2e^{u^2} - 2e^{u^3}; \;\; d_{67}^7 = e^{u^1}; \;\; d_{62}^9 = \tfrac{5}{2}e^{u^1}; \;\; d_{64}^9 = -\tfrac{5}{3}e^{u^4};$$
$$d_{65}^9 = -e^{u^7}; \quad d_{52}^{10} = \tfrac{1}{2}e^{u^1}; \quad d_{34}^{13} = -e^{u^7}; \quad d_{54}^{15} = e^{u^2}.$$

Finally, the matrices G_m, $m = 2, 3, \ldots, 18$, are given by $G_2 = J_7$;

$$G_m = \begin{pmatrix} 1 & 0 \\ 0 & J_6 \end{pmatrix}, \quad m = 3, 4, 5, 6; \quad G_m = \begin{pmatrix} 1 & 0 & 0 \\ 0 & J_5 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad for \;\; m = 7, 8;$$

$$G_m = \begin{pmatrix} E_{5-k} & 0 & 0 \\ 0 & J_k & 0 \\ 0 & 0 & E_2 \end{pmatrix}, \quad where \;\; k = 4 \;\; if \;\; m \; set \; to \; 9, 10;$$

$k = 3$ *if* m *set to* $11, 12$ *and* $k = 2$ *if* m *set to* $13, 14$;

$G_m = E_7, \quad m = 15, 16, 17, 18.$

To find the Laplace invariants of Eqs. (22) when $\alpha = 8$, we introduce matrices of size 8. The matrices P_m, $m = 2, 3, \ldots, 30$, are given by

$$P_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_3 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{2} & 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_4 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{3} & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{6} & \frac{1}{2} & -1 & 1 & 0 & 0 & 0 \\ 0 & \frac{1}{6} & -\frac{1}{2} & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \qquad P_5 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & -\frac{1}{3} & 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{3} & \frac{1}{3} & -1 & 1 & 0 & 0 & 0 \\ 0 & \frac{1}{6} & -\frac{1}{6} & \frac{1}{2} & -1 & 1 & 0 & 0 \\ 0 & -\frac{1}{6} & \frac{1}{6} & -\frac{1}{2} & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_6 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{3}{5} & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{3}{5} & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{5} & \frac{1}{3} & -\frac{1}{3} & 1 & 0 & 0 & 0 \\ 0 & \frac{1}{5} & -\frac{1}{3} & \frac{1}{3} & -1 & 1 & 0 & 0 \\ 0 & -\frac{1}{10} & \frac{1}{6} & -\frac{1}{6} & \frac{1}{2} & -1 & 1 & 0 \\ 0 & \frac{1}{10} & -\frac{1}{6} & \frac{1}{6} & -\frac{1}{2} & 1 & -1 & 1 \end{pmatrix}, \quad P_7 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & \frac{5}{3} & -\frac{5}{3} & 5 & -10 & 10 \\ 0 & 0 & 1 & -\frac{5}{3} & \frac{5}{3} & -5 & 10 & -10 \\ 0 & 0 & 0 & \frac{5}{3} & -\frac{5}{3} & 5 & -10 & 10 \\ 0 & 0 & 0 & 0 & 1 & -3 & 6 & -6 \\ 0 & 0 & 0 & 0 & 0 & 3 & -6 & 6 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix},$$

$$P_8 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{2}{7} & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{2}{7} & 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & \frac{6}{35} & -\frac{3}{5} & \frac{3}{5} & -\frac{3}{5} & 1 & 0 & 0 \\ 0 & -\frac{6}{35} & \frac{3}{5} & -\frac{3}{5} & \frac{3}{5} & -1 & 1 & 0 \\ 0 & \frac{2}{35} & -\frac{1}{5} & \frac{1}{5} & -\frac{1}{5} & \frac{1}{3} & -\frac{1}{3} & 1 \end{pmatrix}, \quad P_9 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{7} & -\frac{12}{35} & \frac{6}{5} & -\frac{6}{5} & \frac{6}{5} & -2 & 0 \\ 0 & -\frac{1}{2} & \frac{6}{5} & -\frac{7}{10} & \frac{7}{10} & -\frac{7}{10} & \frac{7}{6} & 0 \\ 0 & \frac{5}{14} & -\frac{6}{7} & \frac{1}{2} & \frac{1}{2} & -\frac{1}{2} & \frac{5}{6} & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -\frac{5}{3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{5}{3} & 0 \\ 0 & \frac{1}{2} & \frac{1}{5} & -\frac{7}{10} & \frac{7}{10} & -\frac{7}{10} & \frac{7}{6} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_{10} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{7}{6} & \frac{3}{2} & \frac{3}{5} & -\frac{21}{10} & \frac{21}{10} & \frac{1}{15} & 0 \\ 0 & -\frac{1}{3} & \frac{3}{7} & -\frac{8}{35} & \frac{4}{5} & -\frac{4}{5} & \frac{2}{15} & 0 \\ 0 & \frac{7}{6} & -\frac{3}{2} & \frac{4}{5} & \frac{7}{10} & -\frac{7}{10} & -\frac{7}{15} & 0 \\ 0 & -\frac{5}{6} & \frac{15}{14} & -\frac{4}{7} & -\frac{1}{2} & \frac{3}{2} & -\frac{2}{3} & 0 \\ 0 & \frac{7}{6} & -\frac{3}{2} & -\frac{3}{5} & \frac{21}{10} & -\frac{21}{10} & \frac{14}{15} & 0 \\ 0 & \frac{7}{3} & -2 & -\frac{4}{5} & \frac{14}{5} & -\frac{14}{5} & \frac{7}{15} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_{11} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & -1 & \frac{19}{5} & -\frac{19}{5} & 0 & 0 \\ 0 & 0 & 0 & 1 & -\frac{19}{5} & \frac{19}{5} & 0 & 0 \\ 0 & 0 & -1 & \frac{9}{7} & -\frac{24}{35} & \frac{2}{5} & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & \frac{12}{5} & -\frac{7}{5} & 0 & 0 \\ 0 & 0 & 0 & 1 & -\frac{12}{5} & \frac{7}{5} & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \qquad P_{12} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & \frac{7}{2} & -\frac{7}{5} & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\frac{7}{2} & \frac{7}{5} & 1 & 0 & 0 \\ 0 & 0 & 0 & -\frac{7}{2} & 0 & 1 & 0 & 0 \\ 0 & -1 & 1 & 7 & -\frac{7}{5} & -2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_{13} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & \frac{7}{2} & -1 & -1 & 0 & 0 \\ 0 & -1 & 1 & -7 & 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\frac{7}{2} & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & -\frac{7}{2} & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_{14} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & -1 & 1 & -\frac{7}{2} & 2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_{15} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -\frac{1}{2} & -\frac{7}{4} & -\frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{7}{4} & \frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{7} & \frac{1}{2} & \frac{1}{7} & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{7} & -\frac{1}{2} & \frac{1}{7} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_{16} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & \frac{7}{2} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & -\frac{7}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_{17} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & \frac{7}{2} & \frac{7}{2} & 0 & 0 & 0 \\ 0 & 0 & 1 & -\frac{7}{2} & -\frac{7}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_{18} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{2}{7} & -\frac{2}{7} & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_{19} = P_{20} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{2}{7} & -\frac{2}{7} & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_{21} = P_{22} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_{23} = P_{24} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$P_m = 0, \quad m = 25, 26, \ldots, 30.$

The diagonal matrices S_m and R_m, are given by

$$S_1 = \mathrm{diag}\left(\exp(u^1), \exp(u^2), \exp(u^3), \exp(u^4), \exp(u^5), \exp(u^6), \exp(u^7), \exp(u^8)\right),$$
$$S_2 = \mathrm{diag}\left(0, \exp(u^1+u^3), \exp(u^2+u^4), \exp(u^3+u^4), \exp(u^4+u^5), \exp(u^5+u^6), \exp(u^6+u^7),\right.$$
$$\left.\exp(u^7+u^8)\right), \quad S_3 = \mathrm{diag}\left(0, 4\exp(u^2+u^3+u^4), \exp(u^1+u^3+u^4), \exp(u^2+u^4+u^5),\right.$$
$$\left.\exp(u^3+u^4+u^5), \exp(u^4+u^5+u^6), \exp(u^5+u^6+u^7), \exp(u^6+u^7+u^8)\right),$$
$$S_4 = \mathrm{diag}\left(0, 9\exp(u^1+u^2+u^3+u^4), 4\exp(u^2+u^3+u^4+u^5), \exp(u^1+u^3+u^4+u^5),\right.$$
$$\left.\exp(u^2+u^4+u^5+u^6), \exp(u^3+u^4+u^5+u^6), \exp(u^4+u^5+u^6+u^7), \exp(u^5+u^6+u^7+u^8)\right),$$
$$S_5 = \mathrm{diag}\left(0, 4\exp(u^2+u^3+2u^4+u^5), 9\exp(u^1+u^2+u^3+u^4+u^5), 4\exp(u^2+u^3+u^4+u^5+u^6),\right.$$
$$\exp(u^1+u^3+u^4+u^5+u^6), \exp(u^2+u^4+u^5+u^6+u^7), \exp(u^3+u^4+u^5+u^6+u^7),$$
$$\left.\exp(u^4+u^5+u^6+u^7+u^8)\right), \quad S_6 = \mathrm{diag}\left(0, 25\exp(u^1+u^2+u^3+2u^4+u^5),\right.$$
$$4\exp(u^2+u^3+2u^4+u^5+u^6), 9\exp(u^1+u^2+u^3+u^4+u^5+u^6), 4\exp(u^2+u^3+u^4+u^5+u^6+u^7),$$
$$\left.\exp(u^1+u^3+u^4+u^5+u^6+u^7), \exp(u^2+u^4+u^5+u^6+u^7+u^8), \exp(u^3+u^4+u^5+u^6+u^7+u^8)\right),$$
$$S_7 = \mathrm{diag}\left(0, 25\exp(u^1+u^2+2u^3+2u^4+u^5), 4\exp(u^2+u^3+2u^4+2u^5+u^6),\right.$$
$$25\exp(u^1+u^2+u^3+2u^4+u^5+u^6), 4\exp(u^2+u^3+2u^4+u^5+u^6+u^7), 9\exp(u^1+u^2+u^3+u^4+$$
$$\left.+u^5+u^6+u^7), 4\exp(u^2+u^3+u^4+u^5+u^6+u^7+u^8), \exp(u^1+u^3+u^4+u^5+u^6+u^7+u^8), \right),$$
$$S_8 = \mathrm{diag}\left(0, 49\exp(u^1+u^2+u^3+2u^4+2u^5+u^6), 25\exp(u^1+u^2+2u^3+2u^4+u^5+u^6),\right.$$
$$4\exp(u^2+u^3+2u^4+2u^5+u^6+u^7), 25\exp(u^1+u^2+u^3+2u^4+u^5+u^6+u^7),$$
$$\left.4\exp(u^2+u^3+2u^4+u^5+u^6+u^7+u^8), 9\exp(u^1+u^2+u^3+u^4+u^5+u^6+u^7+u^8), 0\right),$$
$$S_9 = \mathrm{diag}\left(0, 4\exp(u^2+u^3+2u^4+2u^5+2u^6+u^7), 49\exp(u^1+u^2+u^3+2u^4+2u^5+u^6+u^7),\right.$$
$$25\exp(u^1+u^2+2u^3+2u^4+u^5+u^6+u^7), 4\exp(u^2+u^3+2u^4+2u^5+u^6+u^7+u^8),$$
$$\left.25\exp(u^1+u^2+u^3+2u^4+u^5+u^6+u^7+u^8), 49\exp(u^1+u^2+2u^3+2u^4+2u^5+u^6), 0\right),$$
$$S_{10} = \mathrm{diag}\left(0, 49\exp(u^1+u^2+2u^3+3u^4+2u^5+u^6), 4\exp(u^2+u^3+2u^4+2u^5+2u^6+u^7+u^8),\right.$$
$$49\exp(u^1+u^2+u^3+2u^4+2u^5+u^6+u^7+u^8), 25\exp(u^1+u^2+2u^3+2u^4+u^5+u^6+u^7+u^8),$$
$$\left.49\exp(u^1+u^2+2u^3+2u^4+2u^5+u^6+u^7), 49\exp(u^1+u^2+u^3+2u^4+2u^5+2u^6+u^7), 0\right),$$

$$S_{11} = \operatorname{diag}\left(0, 49\exp(u^1+2u^2+2u^3+3u^4+2u^5+u^6), 49\exp(u^1+u^2+2u^3+3u^4+2u^5+u^6+u^7),\right.$$
$$4\exp(u^2+u^3+2u^4+2u^5+2u^6+2u^7+u^8), 25\exp(u^1+u^2+2u^3+2u^4+2u^5+u^6+u^7+u^8),$$
$$\left.49\exp(u^1+u^2+u^3+2u^4+2u^5+2u^6+u^7+u^8), 49\exp(u^1+u^2+2u^3+2u^4+2u^5+2u^6+u^7), 0\right),$$

$$S_{12} = \operatorname{diag}\left(0, 49\exp(u^1+2u^2+2u^3+3u^4+2u^5+u^6+u^7), 49\exp(u^1+u^2+2u^3+3u^4+2u^5+\right.$$
$$+2u^6+u^7), 4\exp(u^1+u^2+u^3+2u^4+2u^5+2u^6+2u^7+u^8), 49\exp(u^1+u^2+2u^3+3u^4+2u^5+$$
$$\left.+u^6+u^7+u^8), 49\exp(u^1+u^2+2u^3+2u^4+2u^5+2u^6+u^7+u^8), 0, 0\right),$$

$$S_{13} = \operatorname{diag}\left(0, 49\exp(u^1+2u^2+2u^3+3u^4+2u^5+2u^6+u^7), 49\exp(u^1+u^2+2u^3+3u^4+3u^5+\right.$$
$$+2u^6+u^7), 4\exp(u^1+u^2+2u^3+2u^4+2u^5+2u^6+2u^7+u^8), 49\exp(u^1+2u^2+2u^3+3u^4+$$
$$\left.+2u^5+u^6+u^7+u^8), 49\exp(u^1+u^2+2u^3+3u^4+2u^5+2u^6+u^7+u^8), 0, 0\right),$$

$$S_{14} = \operatorname{diag}\left(0, 49\exp(u^1+2u^2+2u^3+3u^4+3u^5+2u^6+u^7), 49\exp(u^1+u^2+2u^3+3u^4+\right.$$
$$+3u^5+2u^6+u^7+u^8), 4\exp(u^1+u^2+2u^3+3u^4+2u^5+2u^6+2u^7+u^8),$$
$$\left.49\exp(u^1+2u^2+2u^3+3u^4+2u^5+2u^6+u^7+u^8), 0, 0, 0\right),$$

$$S_{15} = \operatorname{diag}\left(0, 49\exp(u^1+2u^2+2u^3+4u^4+3u^5+2u^6+u^7),\right.$$
$$49\exp(u^1+2u^2+2u^3+3u^4+3u^5+2u^6+u^7+u^8), 4\exp(u^1+u^2+2u^3+3u^4+3u^5+2u^6+2u^7+u^8),$$
$$\left.4\exp(u^1+2u^2+2u^3+3u^4+2u^5+2u^6+2u^7+u^8), 0, 0, 0\right),$$

$$S_{16} = \operatorname{diag}\left(0, 49\exp(u^1+2u^2+3u^3+4u^4+3u^5+2u^6+u^7),\right.$$
$$49\exp(u^1+2u^2+2u^3+4u^4+3u^5+2u^6+u^7+u^8), 4\exp(u^1+u^2+2u^3+3u^4+3u^5+3u^6+2u^7+u^8),$$
$$\left.4\exp(u^1+2u^2+2u^3+3u^4+3u^5+2u^6+2u^7+u^8), 0, 0, 0\right),$$

$$S_{17} = \operatorname{diag}\left(0, 49\exp(2u^1+2u^2+3u^3+4u^4+3u^5+2u^6+u^7), 49\exp(u^1+2u^2+3u^3+\right.$$
$$+4u^4+3u^5+2u^6+u^7+u^8), 4\exp(u^1+2u^2+2u^3+3u^4+3u^5+3u^6+2u^7+u^8),$$
$$\left.4\exp(u^1+2u^2+2u^3+4u^4+3u^5+2u^6+2u^7+u^8), 0, 0, 0\right),$$

$$S_{18} = \operatorname{diag}\left(0, 49\exp(2u^1+2u^2+3u^3+4u^4+3u^5+2u^6+u^7+u^8), 49\exp(u^1+2u^2+3u^3+\right.$$
$$\left.+4u^4+3u^5+2u^6+2u^7+u^8), 4\exp(u^1+2u^2+2u^3+4u^4+3u^5+3u^6+2u^7+u^8), 0, 0, 0, 0\right),$$

$$S_{19} = \operatorname{diag}\left(0, 49\exp(2u^1+2u^2+3u^3+4u^4+3u^5+2u^6+2u^7+u^8), 49\exp(u^1+2u^2+\right.$$
$$\left.+3u^3+4u^4+3u^5+3u^6+2u^7+u^8), 4\exp(u^1+2u^2+2u^3+4u^4+4u^5+3u^6+2u^7+u^8), 0, 0, 0, 0\right),$$

$$S_{20} = \operatorname{diag}\left(0, 49\exp(2u^1+2u^2+3u^3+4u^4+3u^5+3u^6+2u^7+u^8),\right.$$
$$\left.49\exp(u^1+2u^2+3u^3+4u^4+4u^5+3u^6+2u^7+u^8), 0, 0, 0, 0, 0\right),$$

$$S_{21} = \operatorname{diag}\left(0, 49\exp(2u^1+2u^2+3u^3+4u^4+4u^5+3u^6+2u^7+u^8), 49\exp(u^1+2u^2+3u^3+\right.$$
$$\left.+5u^4+4u^5+3u^6+2u^7+u^8), 0, 0, 0, 0, 0\right),$$

$$S_{22} = \operatorname{diag}\left(0, 49\exp(2u^1+2u^2+3u^3+5u^4+4u^5+3u^6+2u^7+u^8), 49\exp(u^1+3u^2+3u^3+\right.$$
$$\left.+5u^4+4u^5+3u^6+2u^7+u^8), 0, 0, 0, 0, 0\right),$$

$$S_{23} = \mathrm{diag}\,(0, 49\exp(2u^1 + 2u^2 + 4u^3 + 5u^4 + 4u^5 + 3u^6 + 2u^7 + u^8), 49\exp(2u^1 + 3u^2 +$$

$$+ 3u^3 + 5u^4 + 4u^5 + 3u^6 + 2u^7 + u^8), 0, 0, 0, 0, 0)\,,$$

$$S_{24} = \mathrm{diag}\,(0, 0, 49\exp(2u^1 + 3u^2 + 4u^3 + 5u^4 + 4u^5 + 3u^6 + 2u^7 + u^8), 0, 0, 0, 0, 0)\,,$$

$$S_{25} = \mathrm{diag}\,(0, 0, 49\exp(2u^1 + 3u^2 + 4u^3 + 6u^4 + 4u^5 + 3u^6 + 2u^7 + u^8), 0, 0, 0, 0, 0)\,,$$

$$S_{26} = \mathrm{diag}\,(0, 0, 49\exp(2u^1 + 3u^2 + 4u^3 + 6u^4 + 5u^5 + 3u^6 + 2u^7 + u^8), 0, 0, 0, 0, 0)\,,$$

$$S_{27} = \mathrm{diag}\,(0, 0, 49\exp(2u^1 + 3u^2 + 4u^3 + 6u^4 + 5u^5 + 4u^6 + 2u^7 + u^8), 0, 0, 0, 0, 0)\,,$$

$$S_{28} = \mathrm{diag}\,(0, 0, 49\exp(2u^1 + 3u^2 + 4u^3 + 6u^4 + 5u^5 + 4u^6 + 3u^7 + u^8), 0, 0, 0, 0, 0)\,,$$

$$S_{29} = \mathrm{diag}\,(0, 0, 49\exp(2u^1 + 3u^2 + 4u^3 + 6u^4 + 5u^5 + 4u^6 + 3u^7 + 2u^8), 0, 0, 0, 0, 0)\,, \quad S_{30} = 0,$$

$$R_1 = \mathrm{diag}\left(u^1_y, u^2_y, u^3_y, u^4_y, u^5_y, u^6_y, u^7_y, u^8_y\right),$$

$$R_2 = \mathrm{diag}\left(0, u^1_y + u^3_y, u^2_y + u^4_y, u^3_y + u^4_y, u^4_y + u^5_y, u^5_y + u^6_y, u^6_y + u^7_y, u^7_y + u^8_y\right),$$

$$R_3 = \mathrm{diag}\left(0, u^2_y + u^3_y + u^4_y, u^1_y + u^3_y + u^4_y, u^2_y + u^4_y + u^5_y, u^3_y + u^4_y + u^5_y, u^4_y + u^5_y + u^6_y,\right.$$

$$\left. u^5_y + u^6_y + u^7_y, u^6_y + u^7_y + u^8_y\right),$$

$$R_4 = \mathrm{diag}\left(0, u^1_y + u^2_y + u^3_y + u^4_y, u^2_y + u^3_y + u^4_y + u^5_y, u^1_y + u^3_y + u^4_y + u^5_y,\right.$$

$$\left. u^2_y + u^4_y + u^5_y + u^6_y, u^3_y + u^4_y + u^5_y + u^6_y, u^4_y + u^5_y + u^6_y + u^7_y, u^5_y + u^6_y + u^7_y + u^8_y\right),$$

$$R_5 = \mathrm{diag}\left(0, u^2_y + u^3_y + 2u^4_y + u^5_y, u^1_y + u^2_y + u^3_y + u^4_y + u^5_y,\ u^2_y + u^3_y + u^4_y + u^5_y + u^6_y,\right.$$

$$\left. u^1_y + u^3_y + u^4_y + u^5_y + u^6_y, u^2_y + u^4_y + u^5_y + u^6_y + u^7_y, u^3_y + u^4_y + u^5_y + u^6_y + u^7_y, u^4_y + u^5_y + u^6_y + u^7_y + u^8_y\right),$$

$$R_6 = \mathrm{diag}\left(0, u^1_y + u^2_y + u^3_y + 2u^4_y + u^5_y, u^2_y + u^3_y + 2u^4_y + u^5_y + u^6_y,\ u^1_y + u^2_y + u^3_y + u^4_y + u^5_y + u^6_y,\right.$$

$$u^2_y + u^3_y + u^4_y + u^5_y + u^6_y + u^7_y, u^1_y + u^3_y + u^4_y + u^5_y + u^6_y + u^7_y, u^2_y + u^4_y + u^5_y + u^6_y + u^7_y + u^8_y,$$

$$\left. u^3_y + u^4_y + u^5_y + u^6_y + u^7_y + u^8_y\right),$$

$$R_7 = \mathrm{diag}\left(0, u^1_y + u^2_y + 2u^3_y + 2u^4_y + u^5_y, u^2_y + u^3_y + 2u^4_y + 2u^5_y + u^6_y, u^1_y + u^2_y + u^3_y + 2u^4_y + u^5_y + u^6_y,\right.$$

$$u^2_y + u^3_y + 2u^4_y + u^5_y + u^6_y + u^7_y, u^1_y + u^2_y + u^3_y + u^4_y + u^5_y + u^6_y + u^7_y,$$

$$\left. u^2_y + u^3_y + u^4_y + u^5_y + u^6_y + u^7_y + u^8_y, u^1_y + u^3_y + u^4_y + u^5_y + u^6_y + u^7_y + u^8_y, \right),$$

$$R_8 = \mathrm{diag}\left(0, u^1_y + u^2_y + u^3_y + 2u^4_y + 2u^5_y + u^6_y, u^1_y + u^2_y + 2u^3_y + 2u^4_y + u^5_y + u^6_y,\right.$$

$$u^2_y + u^3_y + 2u^4_y + 2u^5_y + u^6_y + u^7_y, u^1_y + u^2_y + u^3_y + 2u^4_y + u^5_y + u^6_y + u^7_y,$$

$$\left. u^2_y + u^3_y + 2u^4_y + u^5_y + u^6_y + u^7_y + u^8_y, u^1_y + u^2_y + u^3_y + u^4_y + u^5_y + u^6_y + u^7_y + u^8_y, 0\right),$$

$$R_9 = \mathrm{diag}\left(0, u^2_y + u^3_y + 2u^4_y + 2u^5_y + 2u^6_y + u^7_y, u^1_y + u^2_y + u^3_y + 2u^4_y + 2u^5_y + u^6_y + u^7_y,\right.$$

$$u^1_y + u^2_y + 2u^3_y + 2u^4_y + u^5_y + u^6_y + u^7_y, u^2_y + u^3_y + 2u^4_y + 2u^5_y + u^6_y + u^7_y + u^8_y,$$

$$\left. u^1_y + u^2_y + u^3_y + 2u^4_y + + u^5_y + u^6_y + u^7_y + u^8_y, u^1_y + u^2_y + 2u^3_y + 2u^4_y + 2u^5_y + u^6_y, 0\right),$$

$$R_{10} = \mathrm{diag}\left(0, u^1_y + u^2_y + 2u^3_y + 3u^4_y + 2u^5_y + u^6_y, u^2_y + u^3_y + 2u^4_y + 2u^5_y + 2u^6_y + u^7_y + u^8_y,\right.$$

$$u^1_y + u^2_y + u^3_y + 2u^4_y + 2u^5_y + u^6_y + u^7_y + u^8_y, u^1_y + u^2_y + 2u^3_y + 2u^4_y + u^5_y + u^6_y + u^7_y + u^8_y,$$

$$\left. u^1_y + u^2_y + 2u^3_y + 2u^4_y + 2u^5_y + u^6_y + u^7_y, u^1_y + u^2_y + u^3_y + 2u^4_y + 2u^5_y + 2u^6_y + u^7_y, 0\right),$$

$$
\begin{aligned}
&R_{11} = \mathrm{diag}\big(0, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + u_y^6,\ u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + u_y^6 + u_y^7,\\
&u_y^2 + u_y^3 + 2u_y^4 + 2u_y^5 + u_y^6 + 2u_y^7 + u_y^8, u_y^1 + u_y^2 + 2u_y^3 + 2u_y^4 + 2u_y^5 + u_y^6 + u_y^7 + u_y^8,\\
&u_y^1 + u_y^2 + u_y^3 + 2u_y^4 + 2u_y^5 + 2u_y^6 + u_y^7 + u_y^8, u_y^1 + u_y^2 + 2u_y^3 + 2u_y^4 + 2u_y^5 + 2u_y^6 + u_y^7, 0\big)\,,\\
&R_{12} = \mathrm{diag}\big(0, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + u_y^6 + u_y^7, u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + 2u_y^6 + u_y^7,\\
&u_y^1 + u_y^2 + u_y^3 + 2u_y^4 + 2u_y^5 + 2u_y^6 + 2u_y^7 + u_y^8, u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + u_y^6 + u_y^7 + u_y^8,\\
&u_y^1 + u_y^2 + 2u_y^3 + 2u_y^4 + 2u_y^5 + +2u_y^6 + u_y^7 + u_y^8, 0, 0\big)\,,\\
&R_{13} = \mathrm{diag}\big(0, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + 2u_y^6 + u_y^7, u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7,\\
&u_y^1 + u_y^2 + 2u_y^3 + 2u_y^4 + 2u_y^5 + 2u_y^6 + 2u_y^7 + u_y^8, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + u_y^6 + u_y^7 + u_y^8,\\
&u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + 2u_y^6 + u_y^7 + u_y^8, 0, 0\big)\,,\\
&R_{14} = \mathrm{diag}\big(0, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7, u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7 + u_y^8,\\
&u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + 2u_y^6 + 2u_y^7 + u_y^8, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + 2u_y^6 + u_y^7 + u_y^8, 0, 0, 0)\,,\\
&R_{15} = \mathrm{diag}\big(0, u_y^1 + 2u_y^2 + 2u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7,\ u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7 + u_y^8,\\
&u_y^1 + u_y^2 + 2u_y^3 + +3u_y^4 + 3u_y^5 + 2u_y^6 + 2u_y^7 + u_y^8, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 2u_y^5 + 2u_y^6 + 2u_y^7 + u_y^8, 0, 0, 0\big)\,,\\
&R_{16} = \mathrm{diag}\big(0, u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7, u_y^1 + 2u_y^2 + 2u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7 + u_y^8,\\
&u_y^1 + u_y^2 + 2u_y^3 + 3u_y^4 + 3u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8, u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 3u_y^5 + 2u_y^6 + +2u_y^7 + u_y^8, 0, 0, 0\big)\,,\\
&R_{17} = \mathrm{diag}\big(0, 2u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7,\ u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7 + u_y^8,\\
&u_y^1 + 2u_y^2 + 2u_y^3 + 3u_y^4 + 3u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8, u_y^1 + 2u_y^2 + 2u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + 2u_y^7 + u_y^8, 0, 0, 0\big)\,,\\
&R_{18} = \mathrm{diag}\big(0, 2u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + u_y^7 + u_y^8, u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 +\\
&+2u_y^7 + u_y^8, u_y^1 + 2u_y^2 + 2u_y^3 + 4u_y^4 + 3u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8, 0, 0, 0, 0)\,,\\
&R_{19} = \mathrm{diag}\big(0, 2u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 3u_y^5 + 2u_y^6 + 2u_y^7 + u_y^8, u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 3u_y^5 + 3u_y^6 +\\
&+2u_y^7 + u_y^8, u_y^1 + 2u_y^2 + 2u_y^3 + 4u_y^4 + 4u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8, 0, 0, 0, 0)\,,\\
&R_{20} = \mathrm{diag}\big(0, 2u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 3u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8,\ u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 4u_y^5 + 3u_y^6 +\\
&+2u_y^7 + u_y^8, 0, 0, 0, 0, 0)\,,\\
&R_{21} = \mathrm{diag}\big(0, 2u_y^1 + 2u_y^2 + 3u_y^3 + 4u_y^4 + 4u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8,\ u_y^1 + 2u_y^2 + 3u_y^3 + 5u_y^4 + 4u_y^5 + 3u_y^6 +\\
&+2u_y^7 + u_y^8, 0, 0, 0, 0, 0)\,,\\
&R_{22} = \mathrm{diag}\big(0, 2u_y^1 + 2u_y^2 + 3u_y^3 + 5u_y^4 + 4u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8,\ u_y^1 + 3u_y^2 + 3u_y^3 + 5u_y^4 + 4u_y^5 + 3u_y^6 +\\
&+2u_y^7 + u_y^8, 0, 0, 0, 0, 0)\,,\\
&R_{23} = \mathrm{diag}\big(0, 2u_y^1 + 2u_y^2 + 4u_y^3 + 5u_y^4 + 4u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8,\ 2u_y^1 + 3u_y^2 + 3u_y^3 + 5u_y^4 + 4u_y^5 + 3u_y^6 +\\
&+2u_y^7 + u_y^8, 0, 0, 0, 0, 0)\,,\\
&R_{24} = \mathrm{diag}\big(0, 0, 2u_y^1 + 3u_y^2 + 4u_y^3 + 5u_y^4 + 4u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8, 0, 0, 0, 0, 0\big)\,,\\
&R_{25} = \mathrm{diag}\big(0, 0, 2u_y^1 + 3u_y^2 + 4u_y^3 + 6u_y^4 + 4u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8, 0, 0, 0, 0, 0\big)\,,\\
&R_{26} = \mathrm{diag}\big(0, 0, 2u_y^1 + 3u_y^2 + 4u_y^3 + 6u_y^4 + 5u_y^5 + 3u_y^6 + 2u_y^7 + u_y^8, 0, 0, 0, 0, 0\big)\,,
\end{aligned}
$$

$$R_{27} = \operatorname{diag}\left(0, 0, 2u_y^1 + 3u_y^2 + 4u_y^3 + 6u_y^4 + 5u_y^5 + 4u_y^6 + 2u_y^7 + u_y^8, 0, 0, 0, 0, 0\right),$$
$$R_{28} = \operatorname{diag}\left(0, 0, 2u_y^1 + 3u_y^2 + 4u_y^3 + 6u_y^4 + 5u_y^5 + 4u_y^6 + 3u_y^7 + u_y^8, 0, 0, 0, 0, 0\right),$$
$$R_{29} = \operatorname{diag}\left(0, 0, 2u_y^1 + 3u_y^2 + 4u_y^3 + 6u_y^4 + 5u_y^5 + 4u_y^6 + 3u_y^7 + 2u_y^8, 0, 0, 0, 0, 0\right).$$

The matrices Z_m, $m = 1, 2, \ldots, 30$, are given by $Z_1 = U$,

$$Z_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -e^{u^3} & 0 & e^{u^1} & 0 & 0 & 0 & 0 & 0 \\ 0 & -e^{u^4} & 0 & e^{u^2} & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^4} & e^{u^3} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^5} & e^{u^4} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -e^{u^6} & e^{u^5} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -e^{u^7} & e^{u^6} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -e^{u^8} & e^{u^7} \end{pmatrix},$$

$$Z_3 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2e^{u^3} & 2e^{u^2} & 0 & 0 & 0 & 0 \\ 0 & -e^{u^4} & 0 & e^{u^1} & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^5} & 0 & e^{u^2} & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^5} & e^{u^3} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -e^{u^6} & e^{u^4} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -e^{u^7} & e^{u^5} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -e^{u^8} & e^{u^6} \end{pmatrix},$$

$$Z_4 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{3}{2}e^{u^1} & 3e^{u^2} & 0 & 0 & 0 & 0 & 0 \\ 0 & -e^{u^5} & 0 & 2e^{u^3} & 2e^{u^2} & 0 & 0 & 0 \\ 0 & 0 & -e^{u^5} & 0 & e^{u^1} & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^6} & 0 & e^{u^2} & 0 & 0 \\ 0 & 0 & 0 & 0 & -e^{u^6} & e^{u^3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -e^{u^7} & e^{u^4} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -e^{u^8} & e^{u^5} \end{pmatrix},$$

$$Z_5 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^4} & 0 & 0 & 0 & 0 & 0 \\ 0 & -e^{u^5} & \frac{3}{2}e^{u^1} & 3e^{u^2} & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^6} & 0 & 2e^{u^3} & 2e^{u^2} & 0 & 0 \\ 0 & 0 & 0 & -e^{u^6} & 0 & e^{u^1} & 0 & 0 \\ 0 & 0 & 0 & 0 & -e^{u^7} & 0 & e^{u^2} & 0 \\ 0 & 0 & 0 & 0 & 0 & -e^{u^7} & e^{u^3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -e^{u^8} & e^{u^4} \end{pmatrix},$$

$$Z_6 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{5}{2}e^{u^1} & \frac{5}{3}e^{u^4} & 0 & 0 & 0 & 0 & 0 \\ 0 & -e^{u^6} & 0 & e^{u^4} & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^6} & \frac{3}{2}e^{u^1} & 3e^{u^2} & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^7} & 0 & 2e^{u^3} & 2e^{u^2} & 0 \\ 0 & 0 & 0 & 0 & -e^{u^7} & 0 & e^{u^1} & 0 \\ 0 & 0 & 0 & 0 & 0 & -e^{u^8} & 0 & e^{u^2} \\ 0 & 0 & 0 & 0 & 0 & 0 & -e^{u^8} & e^{u^3} \end{pmatrix},$$

$$Z_7 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^3} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^5} & 0 & 0 & 0 & 0 & 0 \\ 0 & -e^{u^6} & \frac{5}{2}e^{u^1} & \frac{5}{3}e^{u^4} & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^7} & 0 & e^{u^4} & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^7} & \frac{3}{2}e^{u^1} & 3e^{u^2} & 0 & 0 \\ 0 & 0 & 0 & 0 & -e^{u^8} & 0 & 2e^{u^3} & 2e^{u^2} \\ 0 & 0 & 0 & 0 & 0 & -e^{u^8} & 0 & e^{u^1} \end{pmatrix},$$

$$Z_8 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{7}{2}e^{u^1} & \frac{7}{5}e^{u^5} & 0 & 0 & 0 & 0 \\ 0 & -e^{u^6} & 0 & e^{u^3} & 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{u^7} & 0 & e^{u^5} & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^7} & \frac{5}{2}e^{u^1} & \frac{5}{3}e^{u^4} & 0 & 0 \\ 0 & 0 & 0 & 0 & -e^{u^8} & 0 & e^{u^4} & 0 \\ 0 & 0 & 0 & 0 & 0 & -e^{u^8} & \frac{3}{2}e^{u^1} & 3e^{u^2} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_9 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & e^{u^6} & 0 & 0 & 0 & 0 \\ 0 & -e^{u^7} & 0 & \frac{7}{2}e^{u^1} & \frac{7}{5}e^{u^5} & 0 & 0 & 0 \\ 0 & 0 & -e^{u^7} & 0 & e^{u^3} & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^8} & 0 & e^{u^5} & 0 & 0 \\ 0 & 0 & 0 & 0 & -e^{u^8} & \frac{5}{2}e^{u^1} & \frac{5}{3}e^{u^4} & 0 \\ 0 & e^{u^3} & \frac{7}{5}e^{u^5} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{10} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & e^{u^4} & 0 \\ 0 & -e^{u^8} & 0 & 0 & e^{u^6} & 0 & 0 & 0 \\ 0 & 0 & -e^{u^8} & 0 & \frac{7}{2}e^{u^1} & \frac{7}{5}e^{u^5} & 0 & 0 \\ 0 & 0 & 0 & -e^{u^8} & 0 & e^{u^3} & 0 & 0 \\ 0 & 0 & -e^{u^3} & -\frac{7}{5}e^{u^5} & 0 & 0 & e^{u^7} & 0 \\ 0 & \frac{7}{2}e^{u^1} & e^{u^6} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{11} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^2} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^7} & 0 & 0 & 0 & e^{u^4} & 0 & 0 \\ 0 & 0 & -e^{u^7} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{5}{7}e^{u^3} & e^{u^5} & \frac{5}{7}e^{u^8} & 0 & 0 \\ 0 & 0 & -\frac{7}{2}e^{u^1} & -e^{u^6} & 0 & 0 & e^{u^8} & 0 \\ 0 & 0 & 0 & 0 & 0 & -e^{u^6} & e^{u^3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{12} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^7} & e^{u^2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^6} & 0 & 0 & 0 & -e^{u^4} & 0 \\ 0 & 0 & 0 & e^{u^1} & 0 & \frac{2}{7}e^{u^7} & 0 & 0 \\ 0 & 0 & e^{u^8} & 0 & \frac{7}{5}e^{u^4} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\frac{7}{5}e^{u^6} & e^{u^3} & e^{u^8} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{13} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^6} & e^{u^2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^5} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & e^{u^3} & 0 & \frac{2}{7}e^{u^7} & 0 & 0 \\ 0 & e^{u^8} & 0 & 0 & e^{u^2} & 0 & 0 & 0 \\ 0 & 0 & -e^{u^8} & 0 & -e^{u^6} & e^{u^4} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{14} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^5} & e^{u^2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^8} & 0 & 0 & -e^{u^5} & 0 & 0 \\ 0 & 0 & 0 & e^{u^4} & 0 & \frac{2}{7}e^{u^7} & 0 & 0 \\ 0 & e^{u^8} & 0 & 0 & e^{u^6} & -e^{u^2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{15} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^4} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^8} & e^{u^2} & 0 & e^{u^5} & 0 & 0 & 0 \\ 0 & 0 & -\frac{2}{7}e^{u^7} & e^{u^5} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^2} & \frac{2}{7}e^{u^7} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{16} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^3} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^8} & e^{u^4} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & e^{u^6} & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{2}{7}e^{u^7} & -e^{u^2} & e^{u^5} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{17} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^1} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^8} & e^{u^3} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -e^{u^2} & e^{u^6} & 0 & 0 & 0 \\ 0 & 0 & \frac{2}{7}e^{u^7} & 0 & e^{u^4} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{18} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^8} & e^{u^1} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^7} & 0 & \frac{7}{2}e^{u^3} & 0 & 0 & 0 \\ 0 & 0 & 0 & e^{u^4} & e^{u^6} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{19} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^7} & e^{u^1} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^6} & \frac{7}{2}e^{u^3} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & e^{u^5} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{20} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^6} & e^{u^1} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^5} & \frac{7}{2}e^{u^3} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{21} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^5} & e^{u^1} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^4} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{22} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^4} & e^{u^1} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{23} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^3} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^2} & e^{u^1} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{24} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{u^2} & e^{u^3} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Z_{m} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{u^{m-21}} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$m = 25, 26, 27, 28, 29, \quad Z_{30} = 0.$

The nonzero elements of the matrices $D_m = (d_{ij})_{i,j=1}^{8}$, $m = 2, 3, \ldots, 30$, are determined by the formulas

$$d_{21}^{3} = -e^{u^3}; \;\; d_{68}^{9} = -e^{u^4}; \;\; d_{78}^{9} = 3e^{u^2} - \tfrac{3}{2}e^{u^1}; \;\; d_{37}^{13} = -e^{u^4}; \;\; d_{67}^{13} = e^{u^8};$$
$$d_{36}^{15} = -e^{u^5}; \quad d_{46}^{15} = \tfrac{2}{7}e^{u^7}; \quad d_{56}^{15} = -e^{u^2}; \quad d_{35}^{19} = \tfrac{7}{2}e^{u^3}, \quad d_{45}^{19} = e^{u^6};$$
$$d_{34}^{21} = \tfrac{7}{2}e^{u^3}; \quad d_{32}^{25} = e^{u^2}.$$

The matrices G_m, $m = 2, 3, \ldots, 30$, are given by

$$G_2 = J_8; \quad G_m = \begin{pmatrix} 1 & 0 \\ 0 & J_7 \end{pmatrix}, \quad m = 3, 4, \ldots, 8;$$

$$G_m = \begin{pmatrix} 1 & 0 & 0 \\ 0 & J_6 & 0 \\ 0 & 0 & 1 \end{pmatrix} \text{ for } \; m = 9, 10, 11; \quad G_m = E_8, \quad m \geq 12.$$

The matrices B_m, $m = 1, 2, \ldots, 29$ of size 8, are defined as

- the first column of the matrices B_m, $m \leq 7$ consist of arbitrary elements,
- for $m = 8, 9, 10$ and 11 in additional to the first column, the last column B_m is also arbitrary,
- the first, seventh and eight columns of the matrices B_m, $m = 12, 13$ consist of arbitrary elements,
- the first, sixth, seventh and eight columns of the matrices B_m, $m = 14, 15, 16, 17$ consist of arbitrary elements,
- the first, firth, sixth, seventh and eight columns of the matrices B_m, $m = 18, 19$ consist of arbitrary elements,
- the first, fourth, firth, sixth, seventh and eight columns of the matrices B_m, $m = 20, 21, 22, 23$ consist of arbitrary elements,
- for $m \geq 24$ in additional to the first, fourth, firth, sixth, seventh and eight columns, the second column B_m is also arbitrary,

with the other element equal to zero.

We note that the values of the index m at which the rank of the generalized invariants X_m drops coincide with the exponents $\begin{cases} 1, 4, 5, 7, 8, 11 \\ 1, 5, 7, 9, 10, 13, 17 \\ 1, 7, 11, 13, 17, 19, 23, 29 \end{cases}$ systems (21) for $\begin{cases} \alpha = 6 \\ \alpha = 7 \\ \alpha = 8 \end{cases}$, corresponding, and the number $\begin{cases} r = 12 \\ r = 18 \\ r = 30 \end{cases}$, for which $X_m = 0$ is equal to the Coxeter number [4].

Acknowledgments. The authors are grateful to V.V. Sokolov for the useful discussions of the results in this work.

This work was supported in part by Russian Foundation for Basic Research (Grant № 05 – 01 – 00775 – a).

References

[1] V.E. Adler and S.Ya. Startsev, *Theor. Math. Phys.*, 121, 1484–1495 (1999).

[2] J.M. Anderson and M. Juras, *Duke. Math. J.*, 89, 351–375 (1997).

[3] J.M. Anderson and N. Kamran, *Duke. Math. J.*, 87, 265–319 (1997).

[4] N. Bourbaki, *Groupes et algébres de Lie* (Chap. 1: Algébres de Lie, Chap. 2: Algébres de Lie libres, Chap. 3: Groupes de Lie; Actualités scientifiques et industrielles 1285, 1349; Éléments de mathématique fascicules 26, 36), Hermann, Paris (1971, 1972).

[5] A.M. Guryeva and A.V. Zhiber, *Theor. Math. Phys.*, 138, 401–421 (2004).

[6] E.V. Ferapontov, *Theor. Math. Phys.*, 110, 68–77 (1997).

[7] A.N. Leznov and M.V. Saveliev, *Group Theory Methods of Integration of Nonlinear Dynamical Systems* [in Russian], Nauka, Moscow (1985).

[8] A.B. Shabat and R.I. Yamilov, "Exponentional systems of type II and Cartan matrices", Preprint №. 1, Bashkir Branch, USSR Acad. Sci., Ufa (1981).

[9] A.B. Shabat, *Phys. Lett. A.*, 200, 121–133 (1995).

[10] V.V. Sokolov and A.V. Zhiber, *Phys. Lett. A.*, 208, 303–308 (1995).

[11] S.Ya. Startsev, *Theor. Math. Phys.*, 127, 460–470 (2001).

[12] S.P. Tsarev, *Theor. Math. Phys.*, 122, 121–133 (2000).

[13] A.V. Zhiber, V.V. Sokolov, and S.Ya. Startsev, *Dokl. Math.*, 52, 128–130 (1995).

[14] A.V. Zhiber and V.V. Sokolov, *Usp. Mat. Nauk*, 56, 61–101 (2001).